

Conformal Yano-Killing tensor for the Kerr metric and conserved quantities

Jacek Jezierski* and Maciej Łukasik

Department of Mathematical Methods in Physics,
University of Warsaw, ul. Hoża 69, 00-681 Warsaw, Poland

PACS numbers: 11.10.Ef, 04.20.Ha, 11.30.Jj

Abstract

Properties of (skew-symmetric) conformal Yano–Killing tensors are reviewed. Explicit forms of three symmetric conformal Killing tensors in Kerr spacetime are obtained from the Yano–Killing tensor. The relation between spin-2 fields and solutions to the Maxwell equations is used in the construction of a new conserved quantity which is quadratic in terms of the Weyl tensor. The formula obtained is similar to the functional obtained from the Bel–Robinson tensor and is examined in Kerr spacetime. A new interpretation of the conserved quantity obtained is proposed.

1 Introduction

According to [28] one can define, in terms of spacetime curvature, two kinds of conserved quantities with the help of conformal Yano–Killing tensors (sometimes called conformal Killing forms or twistor forms, see e.g. [30], [38], [39]). The first kind is linear and the second quadratic with respect to the Weyl tensor but a basis for both of them is the Maxwell field. Conserved quantities which are linear with respect to CYK tensor were investigated many times (cf. [1], [21], [22], [25], [26], [28], [35], [36]). On the other hand, quadratic charges are less known and have usually been examined in terms of the Bel–Robinson tensor (see e.g. [6], [7], [12], [14]).

In [28] the following super-tensor was introduced

$$\mathcal{T}_{\mu\nu\alpha\beta\gamma\delta} = \frac{1}{2} (W_{\mu\sigma\alpha\beta} W_{\nu}{}^{\sigma}{}_{\gamma\delta} + W_{\mu\sigma\gamma\delta} W_{\nu}{}^{\sigma}{}_{\alpha\beta} + W^{*}_{\mu\sigma\alpha\beta} W^{*}_{\nu}{}^{\sigma}{}_{\gamma\delta} + W^{*}_{\mu\sigma\gamma\delta} W^{*}_{\nu}{}^{\sigma}{}_{\alpha\beta}) , \quad (1.1)$$

where $W_{\mu\sigma\alpha\beta}$ is a spin-2 field (Weyl tensor) and by “*” we denote its dual $W^{*}_{\mu\nu\alpha\beta} := \frac{1}{2} W_{\mu\nu\rho\sigma} \varepsilon^{\rho\sigma}{}_{\alpha\beta}$ (cf. Section 3.2 below). The tensor $\mathcal{T}$ has the following algebraic properties:

$$\mathcal{T}_{\mu\nu\alpha\beta\gamma\delta} = \mathcal{T}_{\mu\nu\gamma\delta\alpha\beta} = \mathcal{T}_{(\mu\nu)[\alpha\beta][\gamma\delta]} , \quad \mathcal{T}_{\mu\nu\alpha\beta\gamma\delta} g^{\mu\nu} = 0 , \quad (1.2)$$

*Partially supported by grants KBN 2 P03B 073 24 and EPSRC: EP/D032091/1. E-mail: Jacek.Jezierski@fuw.edu.pl

which are simple consequences of the definition (1.1) and spin-2 field properties (cf. Definition 4 below). A contraction of $\mathcal{T}$ gives the Bel–Robinson tensor:

$$T_{\mu\nu\alpha\gamma}^{BR} = g^{\beta\delta} \mathcal{T}_{\mu\nu\alpha\beta\gamma\delta}.$$

Moreover, in [28] the following properties

$$\nabla^\mu \mathcal{T}_{\mu\nu\alpha\beta\gamma\delta} = 0, \quad \mathcal{T}_{\mu[\nu\alpha\beta]\gamma\delta} = 0 \quad (1.3)$$


of the tensor $\mathcal{T}$ are shown. They enable one to formulate the following

Theorem 1. *If P, Q are CYK tensors, X is a conformal vector field and $\mathcal{T}$ obeys the properties (1.2) and (1.3) then*

$$\nabla^\mu (\mathcal{T}_{\mu\nu\alpha\beta\gamma\delta} X^\nu P^{\alpha\beta} Q^{\gamma\delta}) = 0.$$

The basis of Theorem 1 consists in the observation that the skew-symmetric tensor $F_{\mu\nu} := W_{\mu\nu\alpha\beta} Q^{\alpha\beta}$ fulfils Maxwell’s equations.

We summarize some relationships by the following diagram:



In electrodynamics the linear quantity corresponds to electric or magnetic charge and the quadratic one expresses the energy, linear momentum or angular momentum of the Maxwell field. In gravity both kinds of charges play a role of energy. The linear conserved quantities (as two-surface integrals) correspond to ADM mass and linear

or angular momentum but bilinear ones are not obviously related to energy. They rather play a role of energy estimates like in [12] (cf. [2]). In this paper we propose an interpretation of our quadratic quantity.

We present examples of both kinds of conserved quantities for a stationary rotating black hole described by Kerr spacetime. The linear charge measures (total) ADM mass of black hole in a quasi-local way. On the other hand, the quadratic functional we interpret as a rotational energy of the black hole which is not obvious. Moreover, we derive new conformal (symmetric) Killing tensors which enable one to construct additional constants of motion along null geodesics.

This paper is organized as follows: In the next Section we call some basic facts about CYK two-forms in (pseudo-)Riemannian manifolds. Section 3 contains a small review about CYK tensors and their applications in four-dimensional spacetime. In Section 4 we specify our framework to the case of a stationary rotating black hole described by the Kerr metric. Section 5 is devoted to the conserved quantities which are quadratic in terms of the spin-2 field. One of these charges is applied in Section 6 to the description of rotational energy of the Kerr black hole. To clarify the exposition some of the technical results and proofs have been shifted to the appendix. Moreover, in Appendix B we have placed one theorem about CYK tensors on compact Riemannian manifolds.

2 A short course about CYK tensors

Let M be an n -dimensional ($n > 1$) manifold with a Riemannian or pseudo-Riemannian metric $g_{\mu\nu}$. The covariant derivative associated with the Levi-Civita connection will be denoted by ∇ or just by “;”. By $T_{\dots(\mu\nu)\dots}$ we will denote the symmetric part and by $T_{\dots[\mu\nu]\dots}$ the skew-symmetric part of tensor $T_{\dots\mu\nu\dots}$ with respect to indices μ and ν (analogous symbols will be used for more indices).

Let $Q_{\mu\nu}$ be a skew-symmetric tensor field (two-form) on M and let us denote by $\mathcal{Q}_{\lambda\kappa\sigma}$ a (three-index) tensor which is defined as follows:

$$\mathcal{Q}_{\lambda\kappa\sigma}(Q, g) := Q_{\lambda\kappa;\sigma} + Q_{\sigma\kappa;\lambda} - \frac{2}{n-1} (g_{\sigma\lambda} Q^\nu{}_{\kappa;\nu} + g_{\kappa(\lambda} Q_{\sigma)}{}^\mu{}_{;\mu}) . \quad (2.1)$$

The object $\mathcal{Q}$ has the following algebraic properties

$$\mathcal{Q}_{\lambda\kappa\mu} g^{\lambda\mu} = 0 = \mathcal{Q}_{\lambda\kappa\mu} g^{\lambda\kappa}, \quad \mathcal{Q}_{\lambda\kappa\mu} = \mathcal{Q}_{\mu\kappa\lambda}, \quad (2.2)$$

i.e. it is traceless and partially symmetric.

Definition 1. A skew-symmetric tensor $Q_{\mu\nu}$ is a conformal Yano-Killing tensor (or simply CYK tensor) for the metric g iff $\mathcal{Q}_{\lambda\kappa\sigma}(Q, g) = 0$.

In other words, $Q_{\mu\nu}$ is a conformal Yano-Killing tensor if it fulfils the following equation:

$$Q_{\lambda\kappa;\sigma} + Q_{\sigma\kappa;\lambda} = \frac{2}{n-1} (g_{\sigma\lambda} Q^\nu{}_{\kappa;\nu} + g_{\kappa(\lambda} Q_{\sigma)}{}^\mu{}_{;\mu}) \quad (2.3)$$

(first proposed by Tachibana and Kashiwada, cf. [42]).

A more abstract way with no indices of describing a CYK tensor can be found in [4], [30], [38] or [39], where it is considered as the element of the kernel of the twistor operator $Q \rightarrow \mathcal{T}_{wist}Q$ defined¹ as follows:

$$\forall X \quad \mathcal{T}_{wist}Q(X) := \nabla_X Q - \frac{1}{p+1} X \lrcorner dQ + \frac{1}{n-p+1} g(X) \wedge d^*Q.$$

However, to simplify the exposition, we prefer abstract index notation which also seems to be more popular.

Equation (2.3) may be transformed into the following equivalent form:

$$Q_{\lambda(\kappa;\sigma)} - Q_{\kappa(\lambda;\sigma)} + \frac{3}{n-1} g_{\sigma[\lambda} Q_{\kappa]}{}^\delta{}_{;\delta} = 0 \quad (2.4)$$

and this is a generalization of the equation

$$Q_{\lambda(\kappa;\sigma)} - Q_{\kappa(\lambda;\sigma)} + \eta_{\sigma[\lambda} Q_{\kappa]}{}^\delta{}_{;\delta} = 0, \quad (2.5)$$

which appeared in [36] as the equation for skew-symmetric tensor field $Q_{\mu\nu}$ in Minkowski spacetime with the metric $\eta_{\mu\nu}$. We will show in the sequel that, as in [36], equation (2.3) enables one to use $Q_{\mu\nu}$ for defining conserved charges associated with the Weyl tensor.

Using the following symbol:

$$\xi_\mu := Q^\nu{}_{\mu;\nu}, \quad (2.6)$$

we can rewrite equation (2.3) in the form:

$$Q_{\lambda\kappa;\sigma} + Q_{\sigma\kappa;\lambda} = \frac{2}{n-1} (g_{\sigma\lambda} \xi_\kappa - g_{\kappa(\lambda} \xi_{\sigma)}). \quad (2.7)$$

Let us notice that if $\xi_\mu = 0$, then $Q_{\mu\nu}$ fulfils the equation:

$$Q_{\lambda\kappa;\sigma} + Q_{\sigma\kappa;\lambda} = 0. \quad (2.8)$$

Skew-symmetric tensors fulfilling equation (2.8) are called in the literature Yano tensors (or Yano–Killing tensors; see [3], [18], [40], [44]). It is obvious that a two-form $Q_{\mu\nu}$ is a Yano tensor iff $Q_{\mu\nu;\lambda}$ is totally skew-symmetric in all indices. So, if $Q_{\mu\nu}$ fulfils (2.8), then $\xi_\mu = g^{\kappa\lambda} Q_{\kappa\mu;\lambda} = 0$ (because $g^{\kappa\lambda}$ is symmetric in its indices). That means that each Yano tensor is a conformal Yano–Killing tensor, but not the other

¹Obviously $\mathcal{T}_{wist}Q$ corresponds to tensor $\mathcal{Q}(Q, g)$ (in abstract index notation). Here X is a vector field, Q is a p -form, $g : TM \rightarrow T^*M$ is a Riemannian metric, d^* denotes coderivative etc. Conformal Killing p -forms are defined with the help of natural differential operators on Riemannian manifolds. We know from the representation theory of the orthogonal group, that the space of p -form valued one-forms $(T^*M \otimes \wedge^p T^*M)$ decomposes into the orthogonal and irreducible sum of forms of degree $p+1$ (which gives the exterior differential d), the forms of degree $p-1$ (defined by the coderivative d^*) and the trace-free part of the partial symmetrization (the corresponding first order operator is denoted by $\mathcal{T}_{wist}$).

way around. The necessary and sufficient condition for a CYK tensor to be a Yano tensor is the vanishing of ξ_μ .

CYK tensors are the conformal generalization of Yano tensors. More precisely, for any positive function Ω on M the tensor $\mathcal{Q}$ transforms with respect to the conformal rescaling as follows:

$$\mathcal{Q}_{\lambda\kappa\sigma}(Q, g) = \Omega^{-3} \mathcal{Q}_{\lambda\kappa\sigma}(\Omega^3 Q, \Omega^2 g), \quad (2.9)$$

which implies the following

Theorem 2. *If $Q_{\mu\nu}$ is a CYK tensor for the metric $g_{\mu\nu}$, then $\Omega^3 Q_{\mu\nu}$ is a CYK tensor for the conformally rescaled metric $\Omega^2 g_{\mu\nu}$.*

Proof of the formula (2.9) is given in Appendix A. The form of equation (2.8) does not remain unchanged under conformal rescaling. But that equation is a particular case of equation (2.3) whose form remains unchanged under such a transformation. That means that if Q is a Yano tensor of the metric g , then although in general $\Omega^3 Q$ is not a Yano tensor of the metric $\Omega^2 g$, it is a CYK tensor. In this sense equation (2.3) is a conformal generalization of equation (2.8).

2.1 The connection between CYK tensors and Killing tensors

It is a known fact that the “square” of a Yano tensor is a Killing tensor. It turns out that in the same way CYK tensors are connected with conformal Killing tensors. In the following definitions we will restrict ourselves to the Killing tensors and conformal Killing tensors of rank 2, although one can consider tensors of any rank ([13], [40]).

Definition 2. *A symmetric tensor $A_{\mu\nu}$ is a Killing tensor iff it fulfils the equation:*

$$A_{(\mu\nu;\kappa)} = 0. \quad (2.10)$$

Definition 3. *A symmetric tensor $A_{\mu\nu}$ is a conformal Killing tensor iff it fulfils the equation:*

$$A_{(\mu\nu;\kappa)} = g_{(\mu\nu} A_{\kappa)} \quad (2.11)$$

for a certain covector A_κ .

It is obvious that equation (2.10) is a particular case of (2.11). It is easy to see that the covector A_κ is unambiguously determined by equation (2.11) (it can be shown e.g. by contracting the equation with $g^{\mu\nu}$). To be more precise:

$$A_\kappa = \frac{1}{n+2} (2A^\mu_{\kappa;\mu} + A^\mu_{\mu;\kappa}).$$

From the above definitions one can easily see that a (conformal) Killing tensor is a generalization of a (conformal) Killing vector (cf. [10], [24], [37], [41], [43]).

Obviously, if $Q_{\mu\nu}$ is a skew-symmetric tensor, then $A_{\mu\nu}$ defined by the formula

$$A_{\mu\nu} = Q_{\mu\lambda} Q^\lambda_{\nu} \quad (2.12)$$

is a symmetric tensor. It turns out that if a skew-symmetric tensor $Q_{\mu\nu}$ fulfils equation (2.3) (it is a CYK tensor), then $A_{\mu\nu}$ defined by (2.12) fulfils equation (2.11) with

$$A_\kappa = \frac{2}{n-1} Q_\kappa{}^\lambda Q_\lambda{}^\delta{}_{;\delta} \quad (2.13)$$

(therefore – since $A_{\mu\nu}$ is symmetric – it is a conformal Killing tensor). If $Q_{\mu\nu}$ is a Yano tensor, then A_κ defined by formula (2.13) vanishes, thus $A_{\mu\nu}$ (defined by (2.12)) is a Killing tensor. That enables one to formulate the following (cf. Prop. 5.1. in [18] or 35.44 in [40])

Theorem 3. *If $Q_{\mu\nu}$ and $P_{\mu\nu}$ are (conformal) Yano–Killing tensors, then the symmetrized product $A_{\mu\nu} := Q_{\lambda(\mu} P_{\nu)}{}^\lambda$ is a (conformal) Killing tensor.*

Proof. Let $Q_{\mu\nu}$ and $P_{\mu\nu}$ be conformal Yano-Killing tensors. We have then

$$Q_{\kappa\lambda;\sigma} + Q_{\sigma\lambda;\kappa} = \frac{2}{n-1} (g_{\sigma\kappa} \xi_\lambda - g_{\lambda(\kappa} \xi_{\sigma)})$$

and

$$P_{\kappa\lambda;\sigma} + P_{\sigma\lambda;\kappa} = \frac{2}{n-1} (g_{\sigma\kappa} \zeta_\lambda - g_{\lambda(\kappa} \zeta_{\sigma)}) ,$$

where $\xi_\mu = Q^\nu{}_{\mu;\nu}$ and $\zeta_\mu = P^\nu{}_{\mu;\nu}$. Contracting the first of the above equations with $P_\nu{}^\lambda$, we get

$$Q_{\kappa\lambda;\sigma} P_\nu{}^\lambda + Q_{\sigma\lambda;\kappa} P_\nu{}^\lambda = \frac{2}{n-1} \left(g_{\sigma\kappa} \xi_\lambda P_\nu{}^\lambda - \frac{1}{2} P_{\nu\kappa} \xi_\sigma - \frac{1}{2} P_{\nu\sigma} \xi_\kappa \right) .$$

Symmetrizing this equation in κ, σ and ν , we get (since P is skew-symmetric)

$$Q_{\lambda(\kappa;\sigma} P_{\nu)}{}^\lambda = -\frac{1}{n-1} g_{(\kappa\sigma} P_{\nu)}{}^\lambda \xi_\lambda .$$

Analogously we get

$$Q_{\lambda(\kappa} P_{\nu)}{}^\lambda{}_{;\sigma} = -\frac{1}{n-1} g_{(\kappa\sigma} Q_{\nu)}{}^\lambda \zeta_\lambda .$$

Finally, if $A_{\kappa\nu} := Q_{\lambda(\kappa} P_{\nu)}{}^\lambda$, then

$$\begin{aligned} A_{(\kappa\nu;\sigma)} &= (Q_{\lambda(\kappa} P_{\nu)}{}^\lambda)_{;\sigma} = Q_{\lambda(\kappa;\sigma} P_{\nu)}{}^\lambda + Q_{\lambda(\kappa} P_{\nu)}{}^\lambda{}_{;\sigma} \\ &= -\frac{1}{n-1} (g_{(\kappa\sigma} P_{\nu)}{}^\lambda \xi_\lambda + g_{(\kappa\sigma} Q_{\nu)}{}^\lambda \zeta_\lambda) = g_{(\kappa\nu} A_{\sigma)}, \end{aligned}$$

where

$$A_\nu := \frac{1}{n-1} (P^\lambda{}_\nu \xi_\lambda + Q^\lambda{}_\nu \zeta_\lambda) .$$

This means that $A_{\kappa\nu}$ is a conformal Killing tensor. If now Q and P are Yano tensors, then $\xi_\mu = \zeta_\mu = 0$, which implies $A_\mu = 0$. In that case $A_{\kappa\nu}$ is a Killing tensor. $\square$

2.2 The connection between CYK tensors and Killing vectors

Let us denote by $R^\sigma{}_{\kappa\lambda\mu}$ the Riemann tensor describing the curvature of the manifold (M, g) . We use now an integrability condition

$$2Q_{\lambda\kappa;\nu\mu} = \frac{2}{n-1} (g_{\lambda\mu}\xi_{\kappa;\nu} + g_{\nu\lambda}\xi_{\kappa;\mu} - g_{\mu\nu}\xi_{\kappa;\lambda} - g_{\kappa(\lambda}\xi_{\mu);\nu} + g_{\kappa(\mu}\xi_{\nu);\lambda} - g_{\kappa(\nu}\xi_{\lambda);\mu}) \\ + Q_{\sigma\lambda}R^\sigma{}_{\kappa\mu\nu} + Q_{\sigma\mu}R^\sigma{}_{\kappa\lambda\nu} + Q_{\sigma\nu}R^\sigma{}_{\kappa\lambda\mu} + 2Q_{\sigma\kappa}R^\sigma{}_{\mu\nu\lambda}, \quad (2.14)$$

which we prove in Appendix B.

A contraction in indices κ and ν gives us:

$$g_{\mu\lambda}\xi^\sigma{}_{;\sigma} + (n-2)\xi_{(\mu;\lambda)} = (n-1)R_{\sigma(\mu}Q_{\lambda)}{}^\sigma, \quad (2.15)$$

where by $R_{\mu\nu}$ we denote the Ricci tensor of the metric $g_{\mu\nu}$. Taking the trace of (2.15) we obtain:

$$(2n-2)\xi^\sigma{}_{;\sigma} = (n-1)R_{\sigma\mu}Q^{\mu\sigma} = 0,$$

where the last equality results from the fact that $R_{\sigma\mu}$ is a symmetric tensor and $Q^{\mu\sigma}$ is a skew-symmetric one. Therefore, we have $\xi^\sigma{}_{;\sigma} = 0$ which, for $n > 2$, implies

$$\xi_{(\mu;\lambda)} = \frac{n-1}{n-2}R_{\sigma(\mu}Q_{\lambda)}{}^\sigma. \quad (2.16)$$

If M is an Einstein manifold, i.e. its Ricci tensor $R_{\mu\nu}$ is proportional to its metric $g_{\mu\nu}$, then using equation (2.16) we obtain

$$\xi_{(\mu;\nu)} = -\frac{n-1}{n-2}\lambda g_{\sigma(\mu}Q_{\nu)}{}^\sigma = -\frac{n-1}{n-2}\lambda Q_{(\nu\mu)} = 0.$$

Here $R_{\mu\nu} = -\lambda g_{\mu\nu}$ and by λ we denote a cosmological constant. The condition $\xi_{(\mu;\lambda)} = 0$ means that ξ^μ is a Killing vector field of the metric $g_{\mu\nu}$. That enables one to formulate the following

Theorem 4. *If $g_{\mu\nu}$ is a solution of the vacuum Einstein equations with cosmological constant and $Q_{\mu\nu}$ is its CYK tensor, then $\xi^\mu = \nabla_\nu Q^{\nu\mu}$ is a Killing vector field of the metric $g_{\mu\nu}$.*

Let us notice that this fact reduces the number of Einstein metrics possessing a nontrivial CYK tensor. The existence of a solution of equation (2.3) which is *not* a Yano tensor implies that our manifold M has at least one symmetry. In the case of a Yano tensor that does not have to be true.

3 CYK tensors in four dimensions

In the following Section we restrict ourselves to an oriented manifold M with dimension $n = 4$. It turns out that in this case conformal Yano–Killing tensors possess some additional properties.

3.1 Hodge duality

In the space of differential forms of an oriented manifold one can define a mapping called a Hodge duality (Hodge star). It assigns to every p -form a $(n - p)$ -form (n is the dimension of the manifold). We consider the case of $n = 4$ and $p = 2$. The Hodge star then becomes a mapping which assigns to a two-form ω a two-form $*\omega$. Using coordinates we can express this mapping in the following way:

$$*\omega_{\alpha\beta} := \frac{1}{2}\varepsilon_{\alpha\beta}{}^{\mu\nu}\omega_{\mu\nu}, \quad (3.1)$$

where $\varepsilon_{\alpha\beta\mu\nu}$ is the skew-symmetric Levi-Civita tensor² determining the orientation of the manifold ($\frac{1}{4!}\varepsilon_{\alpha\beta\mu\nu}dx^\alpha \wedge dx^\beta \wedge dx^\mu \wedge dx^\nu$ is the volume form of the manifold M). For Riemannian metrics (with the positive signature) we have $**\omega = \omega$ while for Lorentzian metrics the negative sign appears $**\omega = -\omega$ (we keep assuming that ω is a two-form defined on a four-dimensional manifold M). The last equality shows that in the Lorentzian case we cannot have $*\omega = \pm\omega$, whereas for positive metrics it is possible. If $*\omega = \omega$ ($*\omega = -\omega$), we say that ω is selfdual (anti-selfdual).

Due to a CYK tensor being a two-form, it is reasonable to ask what are the properties of its dual. Let Q be a conformal Yano-Killing tensor and $*Q$ its dual. Moreover, similarly to $\xi_\mu = \nabla^\nu Q_{\nu\mu}$ (cf. (2.6)) let us introduce the following covector

$$\chi_\mu := \nabla^\nu *Q_{\nu\mu}. \quad (3.2)$$

Contracting (3.2) with $\frac{1}{2}\varepsilon_{\alpha\beta}{}^{\lambda\kappa}$ we obtain

$$\nabla_\sigma *Q_{\alpha\beta} = \frac{2}{3}g_{\sigma[\alpha}\chi_{\beta]} + \frac{1}{3}\varepsilon_{\alpha\beta\sigma\kappa}\xi^\kappa. \quad (3.3)$$

If we multiply equation (3.3) by $\frac{1}{2}\varepsilon_{\mu\nu}{}^{\alpha\beta}$, we obtain the analogous equality:

$$Q_{\mu\nu;\sigma} = \frac{2}{3}g_{\sigma[\mu}\xi_{\nu]} \pm \frac{1}{3}\varepsilon_{\mu\nu\sigma\beta}\chi^\beta. \quad (3.4)$$

The plus sign in equation (3.4) refers to metrics with the positive determinant and the minus sign refers to metrics with the negative one.

If we symmetrize equation (3.3) in indices α and σ , we receive the following equality:

$$*Q_{\alpha\beta;\sigma} + *Q_{\sigma\beta;\alpha} = \frac{2}{3}(g_{\sigma\alpha}\chi_\beta - g_{\beta(\alpha}\chi_{\sigma)}) . \quad (3.5)$$

It is not hard to recognize that this is equation (2.3) for the tensor $*Q$. It proves the following

²It can be defined by the formula $\varepsilon_{\alpha\beta\mu\nu} = \sqrt{|\det g|}\epsilon_{\alpha\beta\mu\nu}$, where

$$\epsilon_{\alpha\beta\mu\nu} = \begin{cases} +1 & \text{if } \alpha\beta\mu\nu \text{ is an even permutation of } 0, 1, 2, 3 \\ -1 & \text{if } \alpha\beta\mu\nu \text{ is an odd permutation of } 0, 1, 2, 3 \\ 0 & \text{in any other case} \end{cases}$$

Theorem 5. *Let $g_{\mu\nu}$ be a metric of a four-dimensional differential manifold M . A skew-symmetric tensor $Q_{\mu\nu}$ is a CYK tensor of the metric $g_{\mu\nu}$ iff its dual $*Q_{\mu\nu}$ is a CYK tensor of this metric.*

The above theorem implies that for every four-dimensional manifold solutions of equation (2.3) appear in pairs – to each solution we can assign the solution dual to it (in the Hodge duality sense). Now we can observe that equation (3.4) is equation (3.3), in which Q was replaced by $*Q$ ($\pm$ in equation (3.4) reflects the fact that $**Q = \pm Q$).

Theorems 4 and 5 imply that an Einstein metric for which there exists a solution of equation (2.3) has in general two symmetries. However, it does not have to be so if at least one of the fields ξ^μ or χ^μ is equal to zero (i.e. at least one of the CYK tensors is a Yano tensor) or if those fields are not linearly independent ($\chi^\mu = c\xi^\mu$, $c = \text{const.}$).

3.2 Spin-2 field

In order to be able to use conformal Yano–Killing tensors for the construction of charges in general relativity we have to introduce the notion of a spin-2 field. For the purpose of this paper we introduce the following definition.

Definition 4. *The tensor field $W_{\alpha\beta\mu\nu}$ is called a spin-2 field iff the following equations are fulfilled:*

$$W_{\alpha\beta\mu\nu} = W_{\mu\nu\alpha\beta} = W_{[\alpha\beta][\mu\nu]},$$

$$W_{\alpha[\beta\mu\nu]} = 0, \quad g^{\alpha\mu}W_{\alpha\beta\mu\nu} = 0 \tag{3.6}$$

and

$$\nabla_{[\lambda}W_{\mu\nu]\alpha\beta} = 0. \tag{3.7}$$

According to the above definition the spin-2 field is any Riemann tensor of a vacuum metric (i.e. Ricci flat metric), as well as any Weyl tensor of linearized Einstein theory. Equations (3.7) (which in the case of $W_{\mu\nu\alpha\beta}$ being a Riemann tensor are simply Bianchi identities) can be treated as the field equations for $W_{\mu\nu\alpha\beta}$.

$W_{\mu\nu\alpha\beta}$ is skew-symmetric both in the first and the second pair of indices and to both of them the Hodge star can be applied. We denote

$$*W_{\mu\nu\alpha\beta} = \frac{1}{2}\varepsilon_{\mu\nu}{}^{\rho\sigma}W_{\rho\sigma\alpha\beta}, \quad W^*_{\mu\nu\alpha\beta} = \frac{1}{2}W_{\mu\nu\rho\sigma}\varepsilon^{\rho\sigma}{}_{\alpha\beta}.$$

One can show that

$$*W = W^*, \quad *(*W) = *W^* = \pm W.$$

The “+” sign in the above equation refers to (positive) Riemannian metrics and the “−” sign refers to Lorentzian metrics. One can also show that if W is a spin-2 field, then $*W$ (and consequently W^*) is a spin-2 field as well. If we assume that W fulfils equations (3.6), then the following equivalences are satisfied:

$$\begin{aligned} \nabla_{[\lambda}W_{\mu\nu]\alpha\beta} = 0 &\iff \nabla^\mu W_{\mu\nu\alpha\beta} = 0 \iff \\ \iff \nabla_{[\lambda}*W_{\mu\nu]\alpha\beta} = 0 &\iff \nabla^\mu *W_{\mu\nu\alpha\beta} = 0. \end{aligned} \tag{3.8}$$

3.3 The relation between CYK tensors and spin-2 fields

Let $W_{\mu\nu\alpha\beta}$ be a spin-2 field and $Q_{\mu\nu}$ be any skew-symmetric tensor. Let us denote by F the following two-form

$$F_{\mu\nu}(W, Q) := W_{\mu\nu\lambda\kappa} Q^{\lambda\kappa}. \quad (3.9)$$

The following formula is satisfied:

$$\nabla_\nu F^{\mu\nu}(W, Q) = \frac{2}{3} W^{\mu\nu\alpha\beta} \mathcal{Q}_{\alpha\beta\nu}(Q, g) \quad (3.10)$$

which may be argued as follows:

Proof. Let us notice firstly that due to $W_{\mu\nu\alpha\beta}$ being traceless we have:

$$\begin{aligned} W^{\mu\nu\alpha\beta} \mathcal{Q}_{\alpha\beta\nu} &= W^{\mu\nu\alpha\beta} (Q_{\alpha\beta;\nu} + Q_{\nu\beta;\alpha}) = (W^{\mu\nu\alpha\beta} + W^{\mu\alpha\nu\beta}) Q_{\alpha\beta;\nu} \\ &= (W^{\mu\nu\alpha\beta} + \frac{1}{2} W^{\mu\alpha\nu\beta} - \frac{1}{2} W^{\mu\beta\nu\alpha}) Q_{\alpha\beta;\nu} \\ &= \frac{3}{2} W^{\mu\nu\alpha\beta} Q_{\alpha\beta;\nu}, \end{aligned} \quad (3.11)$$

where we have used the fact that $Q_{\alpha\beta;\nu}$ is skew-symmetric in indices α and β and $W^{\mu[\nu\alpha\beta]} = 0$. Using the above formula (3.11) and properties (3.8) of a spin-2 field $W_{\mu\nu\alpha\beta}$ we obtain the final result

$$\begin{aligned} \nabla_\nu F^{\mu\nu}(W, Q) &= \nabla_\nu (W^{\mu\nu\alpha\beta} Q_{\alpha\beta}) = (\nabla_\nu W^{\mu\nu\alpha\beta}) Q_{\alpha\beta} + W^{\mu\nu\alpha\beta} (\nabla_\nu Q_{\alpha\beta}) \\ &= \frac{2}{3} W^{\mu\nu\alpha\beta} \mathcal{Q}_{\alpha\beta\nu}. \end{aligned}$$

□

Definition 1 means that $\mathcal{Q}_{\alpha\beta\nu}(Q, g) = 0$, hence for a CYK tensor $Q_{\mu\nu}$ we have

$$\nabla_\nu F^{\mu\nu}(W, Q) = 0. \quad (3.12)$$

Let V be a three-volume and ∂V its boundary. Formula (3.12) implies³

$$\int_{\partial V} F^{\mu\nu}(W, Q) d\sigma_{\mu\nu} = \int_V \nabla_\nu F^{\mu\nu}(W, Q) d\Sigma_\mu = 0,$$

which means that the flux of $F^{\mu\nu}$ through any two-dimensional closed surfaces S_1 and S_2 is the same as long as we are able to find a three-volume V between them (i.e. there exists V such that $\partial V = S_1 \cup S_2$). In this sense $Q_{\mu\nu}$ defines a charge related to the spin-2 field⁴ W .

We shall prove now the following

³Symbols $d\sigma_{\mu\nu}$ and $d\Sigma_\mu$ can be defined in the following way: if Ω stands for the volume form of the manifold M , then $d\sigma_{\mu\nu} := (\partial_\mu \wedge \partial_\nu) \lrcorner \Omega$, $d\Sigma_\mu := \partial_\mu \lrcorner \Omega$.

⁴Equation (3.10) can be also used to define asymptotic charges of an asymptotically flat space. Let S_t be the family of closed two-dimensional surfaces going to spatial infinity with $t \rightarrow \infty$. The formula $\lim_{t \rightarrow \infty} \int_{S_t} F^{\mu\nu}(Q, g) d\sigma_{\mu\nu}$ defines a quantity (an asymptotic charge) which does not depend on the choice of the family S_t as long as $\mathcal{Q}_{\mu\nu\rho}(Q, g)$ goes to zero in spatial infinity fast enough (see [22], [25], [28]).

Theorem 6. *Let W be a spin-2 field and Q be a conformal Yano-Killing tensor, then a skew-symmetric tensor $F_{\mu\nu}$ defined by the formula (3.9) satisfies the vacuum Maxwell equations, i.e.*

$$\nabla_\lambda F^{\mu\lambda} = 0 = \nabla_\lambda *F^{\mu\lambda}, \quad (3.13)$$

where $*F^{\mu\lambda} = \frac{1}{2}\varepsilon^{\mu\lambda}{}_{\rho\sigma}F^{\rho\sigma}$.

Proof. The first of equations (3.13) is simply equation (3.10) (since $\mathcal{Q}_{\alpha\beta\nu}(Q, g) = 0$ for Q being a CYK tensor). The second equation is a simple consequence of the fact that $*F(W, Q) = F(*W, Q) = F(W^*, Q) = F(W, *Q)$ and that $*Q$ is a CYK tensor. $\square$

Theorem 6 implies that a charge defined by the surface integral $\int_S F^{\mu\nu} d\sigma_{\mu\nu}$ is the “electromagnetic” charge of the field $F^{\mu\nu}$. More precisely, due to the Levi-Civita connection being torsionless, we have

$$\nabla_{[\lambda} F_{\kappa\rho]} = \partial_{[\lambda} F_{\kappa\rho]}.$$

Moreover,

$$0 = \nabla_\lambda *F^{\mu\lambda} = \nabla_\lambda \left(\frac{1}{2} \varepsilon^{\mu\lambda\kappa\rho} F_{\kappa\rho} \right) = \frac{1}{2} \varepsilon^{\mu\lambda\kappa\rho} F_{\kappa\rho;\lambda} = \frac{1}{2} \varepsilon^{\mu\lambda\kappa\rho} F_{\kappa\rho,\lambda}. \quad (3.14)$$

Multiplying the above equality by $\varepsilon_{\mu\alpha\beta\gamma}$ we get $F_{[\alpha\beta,\gamma]} = 0$, which simply means that $dF = 0$. The formula (3.14) obviously implies that if $dF = 0$, then $\nabla_\lambda *F^{\mu\lambda} = 0$. This means that the equalities (3.13) are equivalent to the following equations:

$$dF = 0 = d *F. \quad (3.15)$$

3.4 CYK tensors in Minkowski spacetime

Let M be the Minkowski space and (x^μ) Cartesian coordinates on M . We have then $g_{\mu\nu} = \eta_{\mu\nu} := \text{diag}(-1, 1, 1, 1)$. Taking into account that the Riemann tensor of the metric $\eta_{\mu\nu}$ vanishes and, in coordinates x^μ , covariant derivative is the same as ordinary derivative (we change “,” into “,”), equation (B.4) takes the following form:

$$Q_{\lambda\kappa,\nu\mu} = \frac{1}{3} \left(\eta_{\lambda\mu} \xi_{\kappa,\nu} + \eta_{\nu\lambda} \xi_{\kappa,\mu} - \eta_{\mu\nu} \xi_{\kappa,\lambda} - \eta_{\kappa(\lambda} \xi_{\mu),\nu} + \eta_{\kappa(\mu} \xi_{\nu),\lambda} - \eta_{\kappa(\nu} \xi_{\lambda),\mu} \right), \quad (3.16)$$

where $Q_{\mu\nu}$ is a CYK tensor on M and $\xi^\nu = \partial_\mu Q^{\mu\nu}$. According to Theorem 4, vanishing of the Ricci tensor implies that ξ^μ is a Killing vector of the metric $\eta_{\mu\nu}$. It is well known that in the Minkowski spacetime the space of Killing fields is spanned by the fields

$$\mathcal{T}_\mu := \frac{\partial}{\partial x^\mu}, \quad \mathcal{L}_{\mu\nu} := x_\mu \frac{\partial}{\partial x^\nu} - x_\nu \frac{\partial}{\partial x^\mu} \quad (3.17)$$

(where $x_\mu = \eta_{\mu\nu} x^\nu$). It means that components of the vector ξ^μ in coordinates (x^μ) are polynomials of degree one in variables x^μ . Equation (3.16) implies that coordinates of the tensor $Q_{\mu\nu}$ also have to be polynomials of degree at most two. We have then $Q_{\mu\nu} = Q^{(0)}_{\mu\nu} + Q^{(1)}_{\mu\nu} + Q^{(2)}_{\mu\nu}$, where $Q^{(i)}_{\mu\nu}$ is a monomial of degree i . If we use this

decomposition in equation (2.3), it becomes obvious that each tensor $Q^{(i)}_{\mu\nu}$ has to fulfil it separately. Let us see what conditions it imposes on those monomials.

First, it is easy to see that every constant skew-symmetric tensor fulfils equation (2.3) and therefore $Q^{(0)}_{\mu\nu}$ stays undetermined. Let us focus then on $Q^{(1)}_{\mu\nu}$. We can write $Q^{(1)}_{\mu\nu} = A_{\mu\nu\sigma}x^\sigma$, where $A_{\mu\nu\sigma}$ is a certain constant tensor (skew-symmetric in indices μ and ν). $Q^{(1)}_{\mu\nu}$ is linear in x^μ and therefore $\xi^{(1)\nu} := Q^{(1)\mu\nu}_{,\mu}$ cannot depend on x^μ . We have then $\xi^{(1)\mu} = u^\mu$, where u^μ denotes a certain constant vector. The equation $Q^{(1)\mu\nu}_{,\mu} = u^\nu$ reduces to $\eta^{\mu\sigma}A_{\mu\nu\sigma} = u_\nu$. Decomposing $A_{\mu\nu\sigma}$ as $A_{\mu\nu\sigma} = \frac{2}{3}\eta_{\sigma[\mu}u_{\nu]} + B_{\mu\nu\sigma}$ we get $B_{\mu\nu\sigma} = B_{[\mu\nu]\sigma}$ and $\eta^{\mu\sigma}B_{\mu\nu\sigma} = 0$. Simple calculation shows that $Q^{(1')}_{\mu\nu} := \frac{2}{3}\eta_{\sigma[\mu}u_{\nu]}x^\sigma = \frac{2}{3}x_{[\mu}u_{\nu]}$ satisfies equation (2.3), which means that tensor $Q^{(1'')}_{\mu\nu} := B_{\mu\nu\sigma}x^\sigma$ also satisfies it. Yet $Q^{(1'')\mu\nu}_{,\mu} := \eta^{\mu\sigma}B_{\mu\nu\sigma} = 0$, which means that $Q^{(1'')\mu\nu}_{,\sigma} + Q^{(1'')\sigma\nu}_{,\mu} = B_{\mu\nu\sigma} + B_{\sigma\nu\mu} = 0$. The last equation is equivalent to the statement that $B_{\mu\nu\sigma} = B_{[\mu\nu]\sigma}$. Since $Q^{(1'')}_{\mu\nu}$ is a CYK tensor linear in x^μ , $*Q^{(1'')}_{\mu\nu}$ also has this property, and what follows $*Q^{(1'')\mu\nu}_{,\mu} = v^\nu$ for a certain constant vector v^ν . The last equation can be written as $\frac{1}{2}\varepsilon^{\sigma\rho\mu\nu}B_{\mu\nu\sigma} = v^\rho$. Multiplying it by $\varepsilon_{\rho\alpha\beta\gamma}$ we get $\varepsilon_{\rho\alpha\beta\gamma}v^\rho = \frac{1}{2}\delta^{\sigma\mu\nu}_{\alpha\beta\gamma}B_{\mu\nu\sigma} = 3B_{[\alpha\beta\gamma]} = 3B_{\alpha\beta\gamma}$. Therefore $Q^{(1'')\mu\nu} = \frac{1}{3}\varepsilon_{\mu\nu\rho\sigma}v^\rho x^\sigma$. Finally $Q^{(1)}_{\mu\nu} = \frac{2}{3}x_{[\mu}u_{\nu]} + \frac{1}{3}\varepsilon_{\mu\nu\rho\sigma}v^\rho x^\sigma$, where u^μ and v^μ are any constant vectors. It is easy to check that every tensor of this form fulfils equation (2.3).

The last thing is computing $Q^{(2)}_{\mu\nu}$. We observe first that the components of the vector $\xi^{(2)\nu} := Q^{(2)\mu\nu}_{,\mu}$ in the coordinates (x^μ) are monomials of degree one in variables x^μ . All Killing fields with this property have the form $k^{\mu\nu}\mathcal{L}_{\mu\nu} = k^{\mu\nu}(x_\mu\partial_\nu - x_\nu\partial_\mu) = 2k_\mu{}^\nu x^\mu\partial_\nu$, where $k^{\mu\nu}$ is a certain constant skew-symmetric tensor. Then let $k^{\mu\nu}$ be such that $\xi^{(2)}_\mu = k_{\mu\nu}x^\nu$. $Q^{(2)}_{\mu\nu}$ can be always written in the form $Q^{(2)}_{\mu\nu} = \frac{1}{2}C_{\mu\nu\alpha\beta}x^\alpha x^\beta$, where $C_{\mu\nu\alpha\beta}$ is a certain constant tensor such that $C_{\mu\nu\alpha\beta} = C_{[\mu\nu](\alpha\beta)}$. It is easy to compute that $Q^{(2)}_{\mu\nu,\sigma} = C_{\mu\nu\sigma\beta}x^\beta$, and it follows that $\xi^{(2)}_\nu = \eta^{\mu\alpha}C_{\mu\nu\alpha\beta}x^\beta = k_{\nu\beta}x^\beta$. Due to the freedom of choice of x^β we can write $\eta^{\mu\alpha}C_{\mu\nu\alpha\beta} = k_{\nu\beta}$. Let us try to decompose $C_{\mu\nu\alpha\beta}$ in a similar way as before: as the sum of "trace" and "traceless" parts. Let

$$C_{\mu\nu\alpha\beta} = a\eta_{\alpha[\mu}k_{\nu]\beta} + a\eta_{\beta[\mu}k_{\nu]\alpha} + b k_{\mu\nu}\eta_{\alpha\beta} + D_{\mu\nu\alpha\beta} , \quad (3.18)$$

where a and b are constants, which we may choose freely, and $D_{\mu\nu\alpha\beta}$ is a certain constant tensor. The above formula and the properties of $C_{\mu\nu\alpha\beta}$ imply that $D_{\mu\nu\alpha\beta} = D_{[\mu\nu](\alpha\beta)}$. We also have

$$k_{\nu\beta} = \eta^{\mu\alpha}C_{\mu\nu\alpha\beta} = (2a - b)k_{\nu\beta} + \eta^{\mu\alpha}D_{\mu\nu\alpha\beta} .$$

Choosing a and b such that $2a - b = 1$, we get $\eta^{\mu\alpha}D_{\mu\nu\alpha\beta} = 0$. If we use the formula $Q^{(2)}_{\mu\nu} = \frac{1}{2}C_{\mu\nu\alpha\beta}x^\alpha x^\beta$ in equation (2.3) then (after using the decomposition (3.18) and ordering of terms) we obtain:

$$D_{\mu\nu\alpha\beta} + D_{\alpha\nu\mu\beta} = \left(\frac{2}{3} - a\right) \left(\eta_{\mu\alpha}k_{\nu\beta} - \frac{1}{2}\eta_{\nu\alpha}k_{\mu\beta} - \frac{1}{2}\eta_{\mu\nu}k_{\alpha\beta} - \frac{3}{2}\eta_{\beta\mu}k_{\nu\alpha} - \frac{3}{2}\eta_{\alpha\beta}k_{\nu\mu} \right) .$$

It is easily seen that if we assume $a = \frac{2}{3}$ (which means $b = 2a - 1 = \frac{1}{3}$), then $D_{\mu\nu\alpha\beta}$ has to satisfy the equation $D_{\mu\nu\alpha\beta} = -D_{\alpha\nu\mu\beta}$. Taking into account that $D_{\mu\nu\alpha\beta} = D_{[\mu\nu](\alpha\beta)}$, we get:

$$D_{\mu\nu\alpha\beta} = -D_{\alpha\nu\mu\beta} = D_{\nu\alpha\beta\mu} = -D_{\beta\alpha\nu\mu} = D_{\alpha\beta\mu\nu} .$$

But that means that $D_{\mu\nu\alpha\beta}$ is at the same time symmetric and skew-symmetric in indices α and β (and in μ and ν as well), and therefore is identically equal to zero. Finally, we get that

$$\begin{aligned} Q^{(2)}_{\mu\nu} &= \left(\frac{1}{3}\eta_{\alpha[\mu}k_{\nu]\beta} + \frac{1}{3}\eta_{\beta[\mu}k_{\nu]\alpha} + \frac{1}{6}k_{\mu\nu}\eta_{\alpha\beta} \right) x^\alpha x^\beta \\ &= \frac{2}{3}x_{[\mu}k_{\nu]\beta}x^\beta + \frac{1}{6}k_{\mu\nu}x_\beta x^\beta. \end{aligned}$$

Summarizing the above considerations, we can write the general form of a solution of equation (2.3) in Minkowski spacetime:

$$Q^{\mu\nu} = q^{\mu\nu} + \frac{2}{3}x^{[\mu}u^{\nu]} + \frac{1}{3}\varepsilon^{\mu\nu}{}_{\kappa\lambda}v^\kappa x^\lambda + \frac{2}{3}x^{[\mu}k^{\nu]\lambda}x_\lambda + \frac{1}{6}k^{\mu\nu}x_\lambda x^\lambda, \quad (3.19)$$

where $q^{\mu\nu}$ and $k^{\mu\nu}$ are any constant skew-symmetric tensors, and u^μ and v^μ are any constant vectors. $\xi^\nu = Q^{\mu\nu}{}_{,\mu}$ for $Q^{\mu\nu}$ given by formula (3.19) is equal to $\xi^\mu = u^\mu + k^{\mu\nu}x_\nu$. Obviously each tensor $q^{\mu\nu}$ and $k^{\mu\nu}$ has six independent components, and each vector u^μ and v^μ has four independent components, which means that the space of solutions of equation (2.3) in Minkowski spacetime is twenty-dimensional⁵.

Let us denote by $\mathcal{D}$ a *dilation vector field*:

$$\mathcal{D} := x^\mu \frac{\partial}{\partial x^\mu}, \quad (3.20)$$

which is involved together with Killing fields $\mathcal{T}_\mu$ and $\mathcal{L}_{\mu\nu}$ (cf. (3.17)) in the following commutation relations:

$$\begin{aligned} [\mathcal{T}_\mu, \mathcal{D}] &= \mathcal{T}_\mu, \quad [\mathcal{D}, \mathcal{L}_{\alpha\beta}] = 0, \\ [\mathcal{T}_\mu, \mathcal{T}_\nu] &= 0, \quad [\mathcal{T}_\mu, \mathcal{L}_{\alpha\beta}] = \eta_{\mu\alpha}\mathcal{T}_\beta - \eta_{\mu\beta}\mathcal{T}_\alpha, \\ [\mathcal{L}_{\mu\nu}, \mathcal{L}_{\alpha\beta}] &= \eta_{\mu\alpha}\mathcal{L}_{\beta\nu} - \eta_{\mu\beta}\mathcal{L}_{\alpha\nu} + \eta_{\nu\alpha}\mathcal{L}_{\mu\beta} - \eta_{\nu\beta}\mathcal{L}_{\mu\alpha}. \end{aligned}$$

Obviously $\mathcal{T}_\mu$ and $\mathcal{L}_{\mu\nu}$ are generators of the Poincaré group. After adding $\mathcal{D}$, we get a set of generators of the pseudo-similarity group (Poincaré group extended by scaling transformation generated by $\mathcal{D}$). Formula (3.19) implies that the following tensors (for $\mu < \nu$) form a basis in the space of solutions of equation (2.5):

$$\mathcal{T}_\mu \wedge \mathcal{T}_\nu, \quad \mathcal{D} \wedge \mathcal{T}_\mu, \quad *(\mathcal{D} \wedge \mathcal{T}_\mu), \quad \mathcal{D} \wedge \mathcal{L}_{\mu\nu} - \frac{1}{2}\eta(\mathcal{D}, \mathcal{D})\mathcal{T}_\mu \wedge \mathcal{T}_\nu. \quad (3.21)$$

Ten of the above CYK tensors, namely $\mathcal{T}_\mu \wedge \mathcal{T}_\nu$ and $*(\mathcal{D} \wedge \mathcal{T}_\mu)$, are Yano tensors.

⁵This is the maximal number of solutions which is achieved by any conformally flat manifold. In general, for the CYK p -forms in an n -dimensional Riemannian manifold M this number equals $\binom{n+2}{p+1}$. Moreover, if a manifold M admits the maximal possible number of linear independent conformal Yano-Killing p -forms, then it is conformally flat.

Using the basis (3.21) we can represent any CYK tensor in Minkowski spacetime as follows:

$$Q = \tilde{q}^{\mu\nu} \mathcal{T}_\mu \wedge \mathcal{T}_\nu + \tilde{u}^\mu \mathcal{D} \wedge \mathcal{T}_\mu + \tilde{v}^\mu *(\mathcal{D} \wedge \mathcal{T}_\mu) + \tilde{k}^{\mu\nu} (\mathcal{D} \wedge \mathcal{L}_{\mu\nu} - \frac{1}{2} \eta(\mathcal{D}, \mathcal{D}) \mathcal{T}_\mu \wedge \mathcal{T}_\nu),$$

where $\tilde{q}^{\mu\nu}$ and $\tilde{k}^{\mu\nu}$ are certain constant skew-symmetric tensors, whereas $\tilde{u}^\mu$ and $\tilde{v}^\mu$ are certain constant vectors.

Let W be a spin-2 field in Minkowski spacetime. We already know that every CYK tensor defines a certain conserved quantity related to the field W . Let us write out the charges given by tensors (3.21). We have:

$$w_{\mu\nu} := \frac{1}{16\pi} \int_{\partial\Sigma} W(\mathcal{T}_\mu \wedge \mathcal{T}_\nu), \quad (3.22)$$

$$p_\mu := \frac{1}{16\pi} \int_{\partial\Sigma} W(\mathcal{D} \wedge \mathcal{T}_\mu), \quad (3.23)$$

$$b_\mu := \frac{1}{16\pi} \int_{\partial\Sigma} *W(\mathcal{D} \wedge \mathcal{T}_\mu), \quad (3.24)$$

$$j_{\mu\nu} := \frac{1}{16\pi} \int_{\partial\Sigma} W \left(\mathcal{D} \wedge \mathcal{L}_{\mu\nu} - \frac{1}{2} \eta(\mathcal{D}, \mathcal{D}) \mathcal{T}_\mu \wedge \mathcal{T}_\nu \right), \quad (3.25)$$

where the notation $W(A \wedge B)$ is to be understood as $W^{\mu\nu}{}_{\alpha\beta} A^\alpha B^\beta d\sigma_{\mu\nu}$. The factor $\frac{1}{16\pi}$ was introduced for normalization.

When W is the Weyl tensor of the linearized Einstein theory, p_μ turns out to be its momentum, b_μ — its dual momentum, and $j_{\mu\nu}$ — its angular momentum tensor⁶ (see [25]). In general, there might also be non-vanishing charges $w_{\mu\nu}$. However, if we assume that $W = O(\frac{1}{r^3})$ (which is a typical situation), then $w_{\mu\nu} = 0$. In this case there are only 14 nontrivial charges but we do not know any local (i.e. using only field equation) argument for this being a general rule. It turns out that vanishing of $w_{\mu\nu}$ is necessary for defining angular momentum of the field at future null infinity (see [22], [28]).

We do not have to restrict ourselves in the above considerations only to the case of Minkowski spacetime. All that what was written above can be easily adapted to the case of asymptotically flat spacetime. By asymptotically flat spacetime we mean a manifold M with a metric g and a set of coordinates (x^μ) such that components of g satisfy the following conditions:

$$g_{\mu\nu} - \eta_{\mu\nu} = O\left(\frac{1}{r}\right), \quad g_{\mu\nu,\kappa} = O\left(\frac{1}{r^2}\right), \quad (3.26)$$

⁶These identifications justify the use of the normalizing factor $\frac{1}{16\pi}$.

where $r := (\sum_{k=1}^3 (x^k)^2)^{\frac{1}{2}}$, and $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$. The coordinates (x^μ) do not have to be global coordinates on M . It is sufficient for them to be defined for r greater than some r_0 (that means that each set of four numbers (x^0, x^1, x^2, x^3) , for which $r > r_0$, defines some point of M).

Now, let W be a spin-2 field defined on M . If the metric admits solutions of equation (2.3), which for r going to infinity go to tensor $\mathcal{D} \wedge \mathcal{T}_\mu$, then we say that p_μ defined by the formula (3.23) defines the μ component of the momentum of the field W . Similarly, we can define dual momentum and angular momentum of the field W , provided that the metric $g_{\mu\nu}$ admits CYK tensors with appropriate asymptotic behavior at spatial infinity.

4 CYK tensor and related objects in Kerr space-time

The Kerr solution to the vacuum Einstein equations is the metric g which in a certain coordinate system (t, r, θ, ϕ) has the form:

$$g = g_{tt} dt^2 + 2g_{t\phi} dt d\phi + g_{rr} dr^2 + g_{\theta\theta} d\theta^2 + g_{\phi\phi} d\phi^2, \quad (4.1)$$

where

$$g_{tt} = -1 + \frac{2mr}{\rho^2}, \quad g_{t\phi} = -\frac{2mra \sin^2 \theta}{\rho^2}, \quad g_{rr} = \frac{\rho^2}{\Delta}, \quad g_{\theta\theta} = \rho^2, \\ g_{\phi\phi} = \sin^2 \theta \left(r^2 + a^2 + \frac{2mra^2 \sin^2 \theta}{\rho^2} \right), \quad (4.2)$$

with

$$\rho^2 = r^2 + a^2 \cos^2 \theta \quad \text{and} \quad \Delta = r^2 - 2mr + a^2. \quad (4.3)$$

This metric describes a rotating black hole of mass m and angular momentum $J = am$. The advantage of the above coordinates (called Boyer–Lindquist coordinates) is that for r much greater than m and a

$$g \approx -dt^2 + dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2,$$

which shows that g approximates the flat metric for r going to infinity. It is not hard to see that equations (3.26) are satisfied, which means that g is an asymptotically flat metric. However, for r equal to $r_\pm := m \pm \sqrt{m^2 - a^2}$ the component g_{rr} is singular (due to $\Delta(r = r_\pm) = 0$). A much better coordinate system for describing the Kerr metric near the surfaces $r = r_\pm$ is Eddington–Finkelstein coordinates: $(v, r, \theta, \tilde{\phi})$, where r and θ are the same as before whereas v and $\tilde{\phi}$ are related to Boyer–Lindquist coordinates via the following formulae:

$$dv = dt + \frac{r^2 + a^2}{\Delta} dr, \quad d\tilde{\phi} = d\phi + \frac{a}{\Delta} dr.$$

In Eddington-Finkelstein coordinates g has the form:

$$g = \tilde{g}_{vv} dv^2 + 2\tilde{g}_{v\tilde{\phi}} dv d\tilde{\phi} + 2\tilde{g}_{r\tilde{\phi}} dr d\tilde{\phi} + 2\tilde{g}_{rv} dr dv + \tilde{g}_{\theta\theta} d\theta^2 + \tilde{g}_{\tilde{\phi}\tilde{\phi}} d\tilde{\phi}^2,$$

where

$$\tilde{g}_{vv} = g_{tt}, \quad \tilde{g}_{v\tilde{\phi}} = g_{t\phi}, \quad \tilde{g}_{r\tilde{\phi}} = -a \sin^2 \theta,$$

$$\tilde{g}_{rv} = 1, \quad \tilde{g}_{\tilde{\phi}\tilde{\phi}} = g_{\phi\phi}, \quad \tilde{g}_{\theta\theta} = g_{\theta\theta}.$$

It can be easily seen that now singularities are absent. Nevertheless, in our considerations we will mostly use Boyer–Lindquist coordinates.

A natural question arises: what are the solutions of equation (2.3) for a rotating black hole described by the Kerr metric? Finding the form of the general CYK tensor in Minkowski spacetime turned out to be quite an easy task, but now the situation is not so simple, as we are dealing with quite a complicated overdetermined system of differential equations for components $Q_{\mu\nu}$. However, there is known in the literature (see [19], [34], [41]) one solution⁷

$$Y := Q_{\text{Kerr}} = r \sin \theta d\theta \wedge [(r^2 + a^2) d\phi - a dt] + a \cos \theta dr \wedge (dt - a \sin^2 \theta d\phi). \quad (4.4)$$

Strictly speaking, it is a solution⁸ of equation (2.8) (i.e. Yano tensor), which – as we know – is a special case of equation (2.3). It means that $\xi^\mu := Y^{\nu\mu}_{;\nu} = 0$.

Since the Kerr metric is asymptotically flat, $Y_{\mu\nu}$ in the neighborhood of spatial infinity should be of the form (3.19). Indeed, asymptotically this tensor looks as follows:

$$Y = r^3 \sin \theta d\theta \wedge d\phi + O(1) = *(\mathcal{T}_0 \wedge \mathcal{D}) + O(1).$$

From (3.24) we see that a charge related to $Y_{\mu\nu}$ is simply $-b_0 = b^0$, which represents dual energy. The Riemann tensor $R_{\mu\nu\rho\sigma}$ of (Ricci flat) Kerr spacetime is simultaneously a Weyl tensor and fulfils properties from Definition 4 of a spin-2 field. Hence we obtain:

$$\begin{aligned} F(R, Y) = & \frac{4ma \sin \theta (r^2 - a^2 \cos^2 \theta)}{\rho^4} dt \wedge d\theta + \frac{8mar \cos \theta}{\rho^4} dt \wedge dr \\ & + \frac{4m \sin \theta (r^2 + a^2) (r^2 - a^2 \cos^2 \theta)}{\rho^4} d\theta \wedge d\phi \\ & + \frac{8ma^2 r \cos \theta \sin^2 \theta}{\rho^4} dr \wedge d\phi, \end{aligned} \quad (4.5)$$

where according to (3.9) $F_{\mu\nu}(R, Y) := R_{\mu\nu\lambda\kappa} Y^{\lambda\kappa}$. We have used symbolic software like MATHEMATICA and WATERLOO MAPLE to check the result (4.5).

⁷According to [24] this is also valid for the charged Kerr solution. Penrose and Walker suggest in [41] than it may be even generalized to the case with cosmological constant. For example, it would be nice to check explicitly existence of CYK tensor for Kerr-Newman-AdS black hole.

⁸To avoid confusion we denote this particular solution by Y .

Direct integration of $F(R, Y)$ on a sphere $r = \text{const.}$ and $t = \text{const.}$ gives

$$\frac{1}{16\pi} \int_S F^{\mu\nu}(R, Y) d\sigma_{\mu\nu} = 0,$$

which means that the dual energy of the Kerr black hole vanishes.

Theorem 5 implies that the dual tensor:

$$*Y = a \cos \theta \sin \theta d\theta \wedge [(r^2 + a^2) d\phi - a dt] + r dr \wedge (a \sin^2 \theta d\phi - dt) \quad (4.6)$$

is also a CYK tensor in Kerr spacetime. The two-form $*Y$ is closed⁹ (i.e. $d*Y = 0$). On the other hand, $dY = 3\rho^2 \sin \theta d\theta \wedge d\phi \wedge dr = 3\partial_t \lrcorner \Omega = 3d\Sigma_t$ (where $\Omega = \rho^2 \sin \theta dt \wedge d\theta \wedge d\phi \wedge dr$ is a volume form) and a vector field $\chi = *Y^{\mu\nu}{}_{;\mu} \partial_\nu = 3\partial_t$ is not vanishing. This implies that $*Y$ is a *Killing potential* which satisfies the CYK equation (cf. [20]).

The asymptotics of tensor $*Y$ look as follows:

$$*Y = r\partial_r \wedge \partial_t + O\left(\frac{1}{r}\right) = \mathcal{D} \wedge \mathcal{T}_0 + O\left(\frac{1}{r}\right).$$

From (3.23) we obtain that the charge related to $*Y$ is p_0 and corresponds to the energy. We have:

$$\begin{aligned} F(R, *Y) &= \frac{8ma^2 r \sin \theta \cos \theta}{\rho^4} d\theta \wedge dt + \frac{4m(r^2 - a^2 \cos^2 \theta)}{\rho^4} dt \wedge dr \\ &\quad + \frac{8mar \sin \theta \cos \theta (r^2 + a^2)}{\rho^4} d\phi \wedge d\theta \\ &\quad + \frac{4ma \sin^2 \theta (r^2 - a^2 \cos^2 \theta)}{\rho^4} dr \wedge d\phi. \end{aligned} \quad (4.7)$$

Integration over a sphere $r = \text{const.}$ and $t = \text{const.}$ gives

$$\frac{1}{16\pi} \int_S *F^{\mu\nu}(R, Y) d\sigma_{\mu\nu} = -m, \quad (4.8)$$

which means that the total energy (mass) of the Kerr black hole equals m . Formula (4.8) expresses mass of a rotating black hole quasi-locally.

Let us denote by A and A' the potentials of the closed two-forms $F(R, Y)$ and $F(R, *Y)$ respectively (cf. 3.15), i.e. $F(R, Y) = dA$ and $F(R, *Y) = dA'$. In Boyer–Lindquist coordinates they look as follows

$$A = \frac{4ma \cos \theta}{\rho^2} dt - \frac{4m \cos \theta (r^2 + a^2)}{\rho^2} d\phi, \quad (4.9)$$

$$A' = \left(\frac{4mr}{\rho^2} - 2 \right) dt - \frac{4mar \sin^2 \theta}{\rho^2} d\phi. \quad (4.10)$$

⁹ Y is a Killing-Yano tensor and corresponding covector ξ vanishes.

The one-forms A and A' are always well defined locally. It turns out that potential A' is global, but A is well defined except at $\theta = 0$ and $\theta = \pi$. Formulae (4.9)–(4.10) are obviously not unique, the potentials A and A' are given up to the gauge $A \rightarrow A + df$ and $A' \rightarrow A' + df'$, where f and f' are arbitrary functions. We would like to stress that A is essentially singular, i.e. all potentials for $F(R, Y)$ have the same singularity¹⁰. The result is not surprising because the two-form $-F(R, *Y)$ describes the “electric” charge m hidden under horizon. Its dual two-form $F(R, Y)$ describes the “magnetic” charge and corresponds to a Dirac monopole. This explains why potential A is not global.

Passing from covectors A and A' to vector fields we obtain

$$g^{-1}(A) = -\frac{4ma \cos \theta}{\rho^2} \partial_t - \frac{4m \cos \theta}{\rho^2 \sin^2 \theta} \partial_\phi \quad (4.11)$$

and

$$g^{-1}(A') = 2\partial_t. \quad (4.12)$$

In particular, $g^{-1}(A')$ is a Killing vector field which implies that $F(R, *Y)$ is a Papapetrou field¹¹.

The Kerr metric is a special case of the electrovac Kerr–Newman metric¹² describing a rotating black hole with electric charge q . It turns out that the two-form $F(R, *Y)$ (given by (4.7)) is related to the Maxwell field $F_{\text{K-N}}$ describing electromagnetic field in Kerr–Newman spacetime. More precisely, both fields have the same form up to the exchange of constants q and $-4m$. For the Kerr–Newman solution the corresponding electromagnetic tensor $F_{\text{K-N}}$ has the following property:

$$F(W, *Y) = -\lim_{q \rightarrow 0} \frac{4m}{q} F_{\text{K-N}}.$$

Finishing this Section let us pass to Eddington–Finkelstein coordinates. Formulae (4.4)–(4.7) and (4.11)–(4.12) take a similar form which corresponds to the change of the names of coordinates from t to v and ϕ to $\tilde{\phi}$ respectively. We have

$$Y = r \sin \theta d\theta \wedge \left[(r^2 + a^2) d\tilde{\phi} - a dv \right] + a \cos \theta dr \wedge \left(dv - a \sin^2 \theta d\tilde{\phi} \right),$$

¹⁰Let us denote by B a gauge equivalent potential to A i.e. $B = A + df$. Integrating A and B along a curve $t = \text{const.}$, $r = \text{const.}$, $\theta = \text{const.}$, $0 < \phi < 2\pi$ we obtain the same result because the function f has to be periodic, $f(\phi = 2\pi) - f(\phi = 0) = 0$. For regular B in the limit $\theta \rightarrow 0$ or $\theta \rightarrow \pi$ our curve goes to a point and the corresponding integral has to vanish. On the other hand, integrating A we obtain $8\pi m \cos \theta (r^2 + a^2) / \rho^2$, which converges to $8\pi m (r^2 + a^2) / \rho^2$ for $\theta \rightarrow 0$ or $\theta \rightarrow \pi$.

¹¹Papapetrou (cf. [15], [31]) pointed out that if K is a Killing vector field for metric g then $F := d(g(K))$ fulfils the Maxwell equations with current $j^\mu := R^\mu{}_\nu K^\nu$, where $R_{\mu\nu}$ is Ricci tensor. In particular, for Ricci flat metric g the two-form F is a solution to vacuum Maxwell equations.

¹²In Boyer–Lindquist coordinates we have

$$g_{\text{K-N}} = -\frac{\Delta}{\rho^2} (dt - a \sin^2 \theta d\phi)^2 + \frac{\sin^2 \theta}{\rho^2} ((r^2 + a^2) d\phi - a dt)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2,$$

where $\Delta = r^2 - 2mr + a^2 + q^2$ and $\rho^2 = r^2 + a^2 \cos^2 \theta$. Moreover, the vector potential $A_{\text{K-N}} = qr (dt - a \sin^2 \theta d\phi) / \rho^2$ is associated with the electromagnetic two-form $F_{\text{K-N}} = dA_{\text{K-N}}$. Obviously, for $q = 0$ we obtain the Kerr metric.

$$*Y = a \cos \theta \sin \theta \, d\theta \wedge \left[(r^2 + a^2) \, d\tilde{\phi} - a \, dv \right] + r \, dr \wedge \left(a \sin^2 \theta \, d\tilde{\phi} - dv \right),$$

etc. Let us notice that the same procedure applied to equations (4.9)–(4.10) leads to

$$\bar{A} = \frac{4ma \cos \theta}{\rho^2} dv - \frac{4m \cos \theta (r^2 + a^2)}{\rho^2} d\tilde{\phi}$$

and

$$\bar{A}' = \left(\frac{4mr}{\rho^2} - 2 \right) dv - \frac{4mar \sin^2 \theta}{\rho^2} d\tilde{\phi},$$

which are not equal to A and A' respectively. However, they are gauge equivalent and describe respectively the same fields $F(R, Y)$ and $F(R, *Y)$.

We denote the conformal Killing tensors associated with Y and $*Y$ by $K_{\mu\nu} := Y_\mu^\kappa Y_{\kappa\nu}$, $K'_{\mu\nu} := *Y_\mu^\kappa *Y_{\kappa\nu}$ and $K''_{\mu\nu} := \frac{1}{2} (Y_\mu^\kappa *Y_{\kappa\nu} + *Y_\mu^\kappa Y_{\kappa\nu})$. Theorem 3 implies that K' and K'' are conformal Killing tensors and, moreover, K is a Killing tensor. In Eddington–Finkelstein coordinates (with raised indices) they look as follows:

$$\begin{aligned} K = & -\frac{a^2 r^2 \sin^2 \theta}{\rho^2} \partial_v \otimes \partial_v - \frac{r^2}{\rho^2} \partial_\theta \otimes \partial_\theta - \frac{r^2}{\rho^2 \sin^2 \theta} \partial_{\tilde{\phi}} \otimes \partial_{\tilde{\phi}} + \frac{a^2 \Delta \cos^2 \theta}{\rho^2} \partial_r \otimes \partial_r \\ & - \frac{2ar^2}{\rho^2} \partial_v \otimes_s \partial_{\tilde{\phi}} + \frac{2a^2(r^2 + a^2) \cos^2 \theta}{\rho^2} \partial_v \otimes_s \partial_r + \frac{2a^3 \cos^2 \theta}{\rho^2} \partial_r \otimes_s \partial_{\tilde{\phi}}, \end{aligned} \quad (4.13)$$

where $a \otimes_s b := \frac{1}{2}(a \otimes b + b \otimes a)$,

$$\begin{aligned} K' = & -\frac{a^4 \sin^2 \theta \cos^2 \theta}{\rho^2} \partial_v \otimes \partial_v - \frac{a^2 \cos^2 \theta}{\rho^2} \partial_\theta \otimes \partial_\theta - \frac{a^2 \cos^2 \theta}{\rho^2 \sin^2 \theta} \partial_{\tilde{\phi}} \otimes \partial_{\tilde{\phi}} + \frac{r^2 \Delta}{\rho^2} \partial_r \otimes \partial_r \\ & - \frac{2a^3 \cos^2 \theta}{\rho^2} \partial_v \otimes_s \partial_{\tilde{\phi}} + \frac{2r^2(r^2 + a^2)}{\rho^2} \partial_v \otimes_s \partial_r + \frac{2ar^2}{\rho^2} \partial_r \otimes_s \partial_{\tilde{\phi}} \end{aligned} \quad (4.14)$$

and

$$\begin{aligned} K'' = & -\frac{ar \cos \theta}{\rho^2} \left(a^2 \sin^2 \theta \partial_v \otimes \partial_v + \partial_\theta \otimes \partial_\theta + \frac{1}{\sin^2 \theta} \partial_{\tilde{\phi}} \otimes \partial_{\tilde{\phi}} + \Delta \partial_r \otimes \partial_r \right. \\ & \left. + 2a \partial_v \otimes_s \partial_{\tilde{\phi}} + 2(r^2 + a^2) \partial_v \otimes_s \partial_r + 2a \partial_r \otimes_s \partial_{\tilde{\phi}} \right). \end{aligned} \quad (4.15)$$

In particular, the traces of our conformal Killing tensors are the following:

$$-\text{tr } K = \text{tr } K' = 2(r^2 - a^2 \cos^2 \theta), \quad \text{tr } K'' = -4ar \cos \theta.$$

The Killing tensor $K^{\mu\nu}$ given in (4.13) is the one which was found in [41] (see also [9], [19], [34]). The constant of motion $K^{\mu\nu} p_\mu p_\nu$ along geodesics, given by

$$g^{\mu\nu} p_\mu \nabla_\nu p = 0, \quad (4.16)$$

is called the Carter constant and it is indispensable for the full description of geodesics in Kerr spacetime (cf. [10], [16], [17], [41]). In this context, our two conformal Killing tensors¹³ K' and K'' produce similar constants of motion along null geodesics. More precisely, (2.11) together with the geodesic equation (4.16) implies that for a null covector p and for any conformal Killing tensor K we have

$$g^{\mu\nu} p_\mu \nabla_\nu (K^{\mu\nu} p_\mu p_\nu) = 0,$$

i.e. $A^{\mu\nu} p_\mu p_\nu$ is a constant of motion along null geodesics.

5 Conserved quantities bilinear in terms of spin-2 field

Let us start with the standard definition of the energy-momentum tensor for a Maxwell field F :

$$T_{\mu\nu}^{\text{EM}}(F) := \frac{1}{8\pi} (F_{\mu\sigma} F_\nu{}^\sigma + F^*_{\mu\sigma} F^{*\nu}{}^\sigma) = \frac{1}{4\pi} \left(F_{\mu\sigma} F_\nu{}^\sigma - \frac{1}{4} g_{\mu\nu} F_{\sigma\rho} F^{\sigma\rho} \right). \quad (5.1)$$

The energy-momentum tensor $T_{\mu\nu}^{\text{EM}}(F)$ is symmetric, traceless and satisfies the following positivity condition: *for any non-spacelike future-directed vector fields X, Y we have $T_{\mu\nu}^{\text{EM}}(F) X^\mu Y^\nu \geq 0$* . Straightforwardly from the definition we get

$$T_{\mu\nu}^{\text{EM}}(F) = T_{\mu\nu}^{\text{EM}}(F^*). \quad (5.2)$$

Moreover, if F is a Maxwell field then

$$\nabla^\mu T_{\mu\nu}^{\text{EM}}(F) = 0, \quad (5.3)$$

and if X is a conformal Killing vector field then the quantity

$$CQ^{EM}(X, \Sigma; F) := \int_\Sigma T_{\mu\nu}^{\text{EM}} X^\mu d\Sigma^\nu$$

defines a global conserved quantity for the spacelike hypersurface Σ with the end at spacelike infinity.

Let us restrict ourselves to the spacelike hyperplanes $\Sigma_t := \{x \in M : x^0 = t = \text{const.}\}$. We use the following convention for indices: (x^μ) $\mu = 0, \dots, 3$ are Cartesian coordinates in Minkowski spacetime, x^0 denotes a temporal coordinate and (x^k) $k = 1, 2, 3$ are coordinates on the spacelike surface Σ_t . If the quantity $CQ^{EM}(X, \Sigma_t)$ is finite for $t = 0$ then it is constant in time. If we want to get a positive definite integral CQ^{EM} , we have to restrict ourselves to the case of a non-spacelike field X . We can choose time translation $\mathcal{T}_0$ or time-like conformal acceleration $\mathcal{K}_0$, where

$$\mathcal{K}_\mu := -2x_\mu \mathcal{D} + x^\sigma x_\sigma \mathcal{T}_\mu \quad (5.4)$$

is a set of four “pure” conformal Killing vector fields which should be added to the eleven fields (3.17) and (3.20) to obtain the full 15-dimensional algebra of the conformal group. This way we get

¹³Unlike the Killing tensor K they seem not to be available in the literature in explicit form.

Theorem 7. *There exist two conserved (time-independent) positive definite integrals $CQ^{EM}(\mathcal{T}_0, \Sigma_t; F)$ and $CQ^{EM}(\mathcal{K}_0, \Sigma_t; F)$ for a field F satisfying the vacuum Maxwell equations.*

Proof. This is a simple consequence of (5.3) and the traceless property of T^{EM} which implies $\nabla^\mu (T_{\mu\nu}^{EM} X^\nu) = 0$ for any conformal Killing vector field X . $\square$

Following [12], for the Bel–Robinson tensor defined as follows:

$$T_{\mu\nu\kappa\lambda}^{BR} := W_{\mu\rho\kappa\sigma} W_{\nu}^{\rho}{}_{\lambda}{}^{\sigma} + W_{\mu\rho\kappa\sigma}^* W_{\nu}^{\rho}{}_{\lambda}{}^{\sigma} \quad (5.5)$$

$$= W_{\mu\rho\kappa\sigma} W_{\nu}^{\rho}{}_{\lambda}{}^{\sigma} + W_{\mu\rho\lambda\sigma} W_{\nu}^{\rho}{}_{\kappa}{}^{\sigma} - \frac{1}{8} g_{\mu\nu} g_{\kappa\lambda} W_{\alpha\beta\gamma\delta} W^{\alpha\beta\gamma\delta}, \quad (5.6)$$

where W is a spin-2 field, one can find a natural generalization of Theorem 7 which is a consequence of the properties similar to (5.3). More precisely, $T_{\mu\nu\kappa\lambda}^{BR}$ is symmetric and traceless in all pairs of indices. Moreover,

$$T^{BR}(W) = T^{BR}(W^*),$$

and if W is a spin-2 field than

$$\nabla^\mu T_{\mu\nu\lambda\kappa}^{BR}(W) = 0. \quad (5.7)$$

If X, Y, Z are conformal Killing vector fields then the quantity

$$CQ^{BR}(X, Y, Z, \Sigma_t; W) := \int_{\Sigma_t} T_{\mu\nu\lambda\kappa}^{BR} X^\mu Y^\nu Z^\lambda d\Sigma^\kappa$$

defines a global charge at time t . The quantity $T^{BR}(X, Y, Z, T)$ is non-negative for any non-spacelike future-directed vector fields X, Y, Z, T whenever at most two of the vector fields are distinct. From the above properties we obtain an extension of Theorem 7 to the case of a spin-2 field W :

Theorem 8. *There exist four conserved (time-independent) positive definite integrals $CQ^{BR}(\mathcal{T}_0, \mathcal{T}_0, \mathcal{T}_0, \Sigma_t; W)$, $CQ^{BR}(\mathcal{K}_0, \mathcal{T}_0, \mathcal{T}_0, \Sigma_t; W)$, $CQ^{BR}(\mathcal{K}_0, \mathcal{K}_0, \mathcal{T}_0, \Sigma_t; W)$ and $CQ^{BR}(\mathcal{K}_0, \mathcal{K}_0, \mathcal{K}_0, \Sigma_t; W)$ for the spin-2 field W satisfying field equations (3.7).*

Proof. Similarly as in Theorem 7, from (5.7) and the traceless property of T^{BR} we get $\nabla^\mu (T_{\mu\nu\lambda\kappa}^{BR} X^\kappa Y^\nu Z^\lambda) = 0$ for any conformal Killing vector fields X, Y, Z . $\square$

In [28] a generalization of the above considerations led to the following functional:

$$\Theta(X, V; Q) := \int_{V \subset \Sigma} T_{\mu\nu}^{EM}(F(W, Q)) X^\mu d\Sigma^\nu \quad (5.8)$$

and according to Theorem 3 in [28], the four conserved quantities CQ^{BR} are contained in the functionals Θ .

In particular, the functional

$$\Theta_0 := \int_V T^{EM}(\mathcal{T}_0, \mathcal{T}_0, F(W, \mathcal{D} \wedge \mathcal{T}_0)) d^3x \quad (5.9)$$

has a positive integrand which is a natural candidate for a certain “energy density”. It contains only first (radial and time) derivatives of quasi-local variables describing linearized gravity. The functional Θ_0 is also very close to the reduced Hamiltonian proposed in [27]. They are proportional to each other after spherical harmonic decomposition (i.e. for each spherical mode).

6 Density of “spin energy” for rotating black hole

The ADM mass of the Kerr black hole (4.2) given by the quasi-local formula (4.8) equals m . On the other hand, the *irreducible mass* of the horizon (see [11], [17]) is defined by the quadratic relation to its two-dimensional area A_H :

$$M_{irr} = \sqrt{\frac{A_H}{16\pi}} = \frac{1}{2} \sqrt{r_+^2 + a^2} = \frac{1}{2} \sqrt{2mr_+}.$$

The energy M_{irr} plays the role of a lower bound in Penrose inequality (cf. [8], [33]) and never decreases according to the second law of black hole physics (see e.g. [17] or [23]). The difference between total ADM mass m and the irreducible mass M_{irr} is often interpreted as *rotational energy*

$$M_{rot} := m - M_{irr} = m \left(1 - \sqrt{\frac{r_+}{2m}} \right) \approx a^2/8m.$$

Contrary to M_{irr} the energy M_{rot} is accessible by reversible transformations like the Penrose process [33] which takes place in the (ergo-)region¹⁴ between event horizon and ergo-surface¹⁵.

The two-form (4.7), previously used in Section 4 to obtain the quasi-local mass, may be also applied to the construction of a quadratic functional (5.9) denoted by Θ_0 . Let us consider an auxiliary Maxwell field f which is a solution of the vacuum Maxwell equations on the Kerr background and is given by

$$\begin{aligned} f_{aux} = F(R, *Y)/4m = & \frac{2a^2 r \sin \theta \cos \theta}{\rho^4} d\theta \wedge dt + \frac{r^2 - a^2 \cos^2 \theta}{\rho^4} dt \wedge dr \\ & + \frac{2ar \sin \theta \cos \theta (r^2 + a^2)}{\rho^4} d\phi \wedge d\theta \\ & + \frac{a \sin^2 \theta (r^2 - a^2 \cos^2 \theta)}{\rho^4} dr \wedge d\phi. \end{aligned} \quad (6.1)$$

The density of the energy built from f and corresponding to the integrand in the functional Θ_0 takes the following form

$$\varepsilon_{EM}(f) = \sqrt{-g} T^0_0(f) = \frac{\sqrt{-g}}{8\pi} (E^2 + B^2) = \frac{\sqrt{-g^{00}}}{8\pi} \left(g^{kl} E_k E_l + \frac{1}{2} B_{kl} B_{mn} g^{km} g^{ln} \right),$$

¹⁴Usually called *ergosphere* which is completely misleading because it is not a sphere.

¹⁵It is given by the largest root of $\Delta(r) = a^2 \sin^2 \theta$, usually called the *stationary limit surface*. Moreover, the instantaneous ergo-surface is a two-dimensional manifold with spherical topology and conical singularity on the axis ($\theta = 0, \pi$) (cf. [32]).

where the energy-momentum tensor T is given by (5.1), $E_k = f_{0k}$ and $B_{kl} = f_{kl}$. Formula (5.1) implies that the tensor f and its dual field

$$\begin{aligned} *f_{aux} := F(R, Y)/4m = & \frac{a \sin \theta (r^2 - a^2 \cos^2 \theta)}{\rho^4} dt \wedge d\theta + \frac{2ar \cos \theta}{\rho^4} dt \wedge dr \\ & + \frac{\sin \theta (r^2 + a^2) (r^2 - a^2 \cos^2 \theta)}{\rho^4} d\theta \wedge d\phi + \\ & + \frac{2a^2 r \cos \theta \sin^2 \theta}{\rho^4} dr \wedge d\phi \end{aligned} \quad (6.2)$$

give the same density ε_{EM} .

For f_{aux} given by (6.1) we get the following non-vanishing components:

$$\begin{aligned} E_r = \frac{r^2 - a^2 \cos^2 \theta}{\rho^4}, \quad E_\theta = -\frac{2a^2 r \sin \theta \cos \theta}{\rho^4}, \\ B_{r\phi} = \frac{a \sin^2 \theta (r^2 - a^2 \cos^2 \theta)}{\rho^4}, \quad B_{\theta\phi} = -\frac{2ar \sin \theta \cos \theta (r^2 + a^2)}{\rho^4}, \end{aligned}$$

and finally

$$\varepsilon_{EM}(f) = \frac{\sin \theta (r^2 + a^2)^2 + a^2 \Delta \sin^2 \theta}{8\pi \rho^2 (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta}. \quad (6.3)$$

Our aim is to show that the “spin energy density” coming from functional (5.9) is just the rescaled density (6.3). For this purpose let us consider another Maxwell field (proportional to f)

$$F_{rot} = C(\alpha) \frac{a}{\sqrt{2}} f_{aux},$$

where $\alpha := a/m$ and the rescaling function $C(\alpha)$ will be specified in the sequel. Obviously (6.3) gives

$$\varepsilon_{rot}(F) = \frac{C^2 a^2 \sin \theta (r^2 + a^2)^2 + a^2 \Delta \sin^2 \theta}{16\pi \rho^2 (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta}. \quad (6.4)$$

Integrating density $\varepsilon_{rot}(F)$ outside of the horizon we obtain a total “spin energy”:

$$M_{rot} = 2\pi \int_{r_+}^{\infty} dr \int_0^\pi d\theta \varepsilon_{rot}. \quad (6.5)$$

Using in the integral (6.5) new variables $y = r/m$ and $x = \cos \theta$ it is easy to rewrite it in the form

$$M_{rot} = \frac{C^2 a^2}{8m} \Psi(\alpha),$$

where

$$\Psi(\alpha) := \int_{1+\sqrt{1-\alpha^2}}^{\infty} dy \int_{-1}^1 \frac{dx}{y^2 + \alpha^2 x^2} \frac{(y^2 + \alpha^2)^2 + \alpha^2 (y^2 - 2y + \alpha^2)(1 - x^2)}{(y^2 + \alpha^2)^2 - \alpha^2 (y^2 - 2y + \alpha^2)(1 - x^2)}. \quad (6.6)$$

The “constant” C is defined by the relation which expresses a balance of the energy:

$$m = M_{irr} + M_{rot}. \quad (6.7)$$

Hence

$$C^2 = \frac{8 \left(1 - \sqrt{\frac{1+\sqrt{1-\alpha^2}}{2}} \right)}{\alpha^2 \Psi}. \quad (6.8)$$

The integral (6.6), which is not elementary, may be easily approximated with the help of symbolic software like MATHEMATICA or WATERLOO MAPLE and it gives as follows

$$\begin{aligned} \Psi(\alpha) &= 1 + \frac{\alpha^2}{4} + \frac{33\alpha^4}{280} + \frac{3059\alpha^6}{43200} + \frac{835171\alpha^8}{17297280} + O(\alpha^9), \\ \frac{8 \left(1 - \sqrt{\frac{1+\sqrt{1-\alpha^2}}{2}} \right)}{\alpha^2} &= 1 + \frac{5\alpha^2}{16} + \frac{21\alpha^4}{128} + \frac{429\alpha^6}{4096} + \frac{2431\alpha^8}{32768} + O(\alpha^{10}) \end{aligned}$$

and

$$C = 1 + \frac{\alpha^2}{32} + \frac{1061\alpha^4}{71680} + \frac{397907\alpha^6}{44236800} + \frac{245297388031\alpha^8}{39675808972800} + O(\alpha^9).$$

However, for $\alpha = 0$ the integral (6.6) is elementary and gives $\Psi(0) = 1 = C(0)$. Formula (6.4) together with (6.8) defines uniquely the “spin energy density” coming from (5.9) and compatible with (6.7).

A Conformal rescaling of CYK tensors

In this section we will be dealing with conformal transformations and their impact on conformal Yano-Killing tensors. Since most of the considerations here are independent on the dimension of a manifold, we will not restrict ourselves to spacetime of dimension four. We assume that we are dealing with n -dimensional manifold and that this manifold has a metric g (signature of g plays no role).

Let $\Gamma^\alpha_{\mu\nu}$ denotes Christoffel symbols of Levi-Civita connection associated with the metric g . We have:

$$\Gamma^\alpha_{\mu\nu} = \frac{1}{2} g^{\alpha\beta} (g_{\beta\mu,\nu} + g_{\beta\nu,\mu} - g_{\mu\nu,\beta}). \quad (A.1)$$

Let $\tilde{g}$ be conformally rescaled metric, i.e. $\tilde{g}_{\mu\nu} := \Omega^2 g_{\mu\nu}$ (and what follows, $\tilde{g}^{\mu\nu} := \Omega^{-2} g^{\mu\nu}$), where Ω is a certain positive function ($\Omega > 0$). We will denote Christoffel symbols of this metric by $\tilde{\Gamma}^\alpha_{\mu\nu}$, and covariant derivative associated with them by $\tilde{\nabla}_\mu$.

Obviously, for $\tilde{g}_{\mu\nu}$ and $\tilde{\Gamma}^\alpha_{\mu\nu}$ we have

$$\begin{aligned}
\tilde{\Gamma}^\alpha_{\mu\nu} &= \frac{1}{2}\tilde{g}^{\alpha\beta}(\tilde{g}_{\beta\mu,\nu} + \tilde{g}_{\beta\nu,\mu} - \tilde{g}_{\mu\nu,\beta}) \\
&= \frac{1}{2}\Omega^{-2}g^{\alpha\beta}((\Omega^2 g_{\beta\mu})_{,\nu} + (\Omega^2 g_{\beta\nu})_{,\mu} + (\Omega^2 g_{\mu\nu})_{,\beta}) \\
&= \Gamma^\alpha_{\mu\nu} + \frac{1}{2}\Omega^{-2}g^{\alpha\beta}((\Omega^2)_{,\nu}g_{\beta\mu} + (\Omega^2)_{,\mu}g_{\beta\nu} + (\Omega^2)_{,\beta}g_{\mu\nu}) \\
&= \Gamma^\alpha_{\mu\nu} + g^{\alpha\beta}(U_{,\nu}g_{\beta\mu} + U_{,\mu}g_{\beta\nu} + U_{,\beta}g_{\mu\nu}) \\
&= \Gamma^\alpha_{\mu\nu} + \delta^\alpha_\mu U_{,\nu} + \delta^\alpha_\nu U_{,\mu} - g^{\alpha\beta}U_{,\beta}g_{\mu\nu}, \tag{A.2}
\end{aligned}$$

which is analogous to (A.1). Here $U := \log \Omega$. Using the formula (A.2) and the following identities:

$$\nabla_\mu X_{\nu\rho} = X_{\nu\rho,\mu} - X_{\alpha\rho}\Gamma^\alpha_{\nu\mu} - X_{\nu\alpha}\Gamma^\alpha_{\rho\mu}, \tag{A.3}$$

$$\tilde{\nabla}_\mu X_{\nu\rho} = X_{\nu\rho,\mu} - X_{\alpha\rho}\tilde{\Gamma}^\alpha_{\nu\mu} - X_{\nu\alpha}\tilde{\Gamma}^\alpha_{\rho\mu} \tag{A.4}$$

(which are true for any tensor $X_{\mu\nu}$), we get:

$$\begin{aligned}
\tilde{\nabla}_\mu X_{\nu\rho} &= \nabla_\mu X_{\nu\rho} - X_{\mu\rho}U_{,\nu} - X_{\nu\mu}U_{,\rho} - 2X_{\nu\rho}U_{,\mu} + \\
&\quad + g^{\alpha\beta}U_{,\beta}(X_{\alpha\rho}g_{\mu\nu} + X_{\nu\alpha}g_{\mu\rho}). \tag{A.5}
\end{aligned}$$

We will now prove the formula (2.9). Let us introduce the following notation:

$$\tilde{Q}_{\mu\nu} = \Omega^3 Q_{\mu\nu}, \tag{A.6}$$

$$\tilde{\xi}_\rho := \tilde{g}^{\mu\nu}\tilde{\nabla}_\mu \tilde{Q}_{\nu\rho}. \tag{A.7}$$

Using the formula (A.5), we can rewrite the formula (A.7) in the following form:

$$\begin{aligned}
\tilde{\xi}_\rho &= \tilde{g}^{\mu\nu}\nabla_\mu \tilde{Q}_{\nu\rho} - 3\tilde{g}^{\mu\nu}\tilde{Q}_{\mu\rho}U_{,\nu} + \Omega^{-2}g^{\mu\nu}g^{\alpha\beta}U_{,\beta}(\tilde{Q}_{\alpha\rho}g_{\mu\nu} + \tilde{Q}_{\nu\alpha}g_{\mu\rho}) \\
&= \tilde{g}^{\mu\nu}\nabla_\mu \tilde{Q}_{\nu\rho} - 3\tilde{g}^{\mu\nu}\tilde{Q}_{\mu\rho}U_{,\nu} + \tilde{g}^{\alpha\beta}U_{,\beta}(n\tilde{Q}_{\alpha\rho} + \tilde{Q}_{\nu\alpha}\delta^\nu_\rho) \\
&= \tilde{g}^{\mu\nu}\nabla_\mu \tilde{Q}_{\nu\rho} + (n-4)\tilde{g}^{\mu\nu}\tilde{Q}_{\mu\rho}U_{,\nu} \\
&= \Omega^{-2}g^{\mu\nu}\nabla_\mu(\Omega^3 Q_{\nu\rho}) + (n-4)\Omega g^{\mu\nu}Q_{\mu\rho}U_{,\nu} \\
&= \Omega g^{\mu\nu}\nabla_\mu Q_{\nu\rho} + \Omega^{-2}g^{\mu\nu}(\Omega^3)_{,\mu}Q_{\nu\rho} + (n-4)\Omega g^{\mu\nu}Q_{\mu\rho}U_{,\nu} \\
&= \Omega(\xi_\rho + (n-1)g^{\mu\nu}Q_{\mu\rho}U_{,\nu}), \tag{A.8}
\end{aligned}$$

where $\xi_\rho := g^{\mu\nu} \nabla_\mu Q_{\nu\rho}$. The last equality in (A.8) is a result of the following identities:
 $\Omega^{-2}(\Omega^3)_{,\mu} = 3\Omega_{,\mu} = 3\Omega(\log \Omega)_{,\mu} = 3\Omega U_{,\mu}$. Using the formula (A.5) for $\tilde{Q}$ we get:

$$\begin{aligned} \tilde{\nabla}_\mu \tilde{Q}_{\nu\rho} = \Omega^3 & \left[\nabla_\mu Q_{\nu\rho} + Q_{\nu\rho} U_{,\mu} - Q_{\nu\mu} U_{,\rho} - Q_{\mu\rho} U_{,\nu} \right. \\ & \left. + g^{\alpha\beta} U_{,\beta} (g_{\mu\nu} Q_{\alpha\rho} + g_{\mu\rho} Q_{\nu\alpha}) \right] \end{aligned} \quad (\text{A.9})$$

and what follows:

$$\begin{aligned} \tilde{\nabla}_\mu \tilde{Q}_{\nu\rho} + \tilde{\nabla}_\rho \tilde{Q}_{\nu\mu} = \Omega^3 & \left[\nabla_\mu Q_{\nu\rho} + \nabla_\rho Q_{\nu\mu} + g^{\alpha\beta} U_{,\beta} (g_{\mu\nu} Q_{\alpha\rho} + \right. \\ & \left. + g_{\rho\nu} Q_{\alpha\mu} + 2g_{\mu\rho} Q_{\nu\alpha}) \right]. \end{aligned} \quad (\text{A.10})$$

Formulae (2.1), (A.8) and (A.10) imply that

$$\begin{aligned} \mathcal{Q}_{\mu\nu\rho}(\tilde{Q}, \tilde{g}) &= \tilde{\nabla}_\rho \tilde{Q}_{\mu\nu} + \tilde{\nabla}_\mu \tilde{Q}_{\rho\nu} - \frac{2}{n-1} \left(\tilde{g}_{\mu\rho} \tilde{\xi}_\nu - \tilde{g}_{\nu(\mu} \tilde{\xi}_{\rho)} \right) \\ &= \Omega^3 \left[\nabla_\rho Q_{\mu\nu} + \nabla_\mu Q_{\rho\nu} - \frac{2}{n-1} (g_{\mu\rho} \xi_\nu - g_{\nu(\mu} \xi_{\rho)}) \right] \\ &= \Omega^3 \mathcal{Q}_{\mu\nu\rho}(Q, g), \end{aligned} \quad (\text{A.11})$$

which proves the formula (2.9).

B Integrability conditions for the CYK equation

For a manifold M equipped with metric tensor $g_{\mu\nu}$ and for any tensor field $Q_{\mu\nu}$ on M the following equality holds

$$Q_{\lambda\kappa;\nu\mu} - Q_{\lambda\kappa;\mu\nu} = Q_{\sigma\kappa} R^\sigma{}_{\lambda\nu\mu} + Q_{\lambda\sigma} R^\sigma{}_{\kappa\nu\mu}, \quad (\text{B.1})$$

where $R^\mu{}_{\nu\rho\sigma}$ is Riemann tensor of metric $g_{\mu\nu}$. Obviously we assume that Q is skew-symmetric which implies that

$$2\xi^\mu{}_{;\mu} = Q^{\lambda\kappa}{}_{;\lambda\kappa} - Q^{\lambda\kappa}{}_{;\kappa\lambda} = 2Q^{\kappa\sigma} R_{\sigma\kappa} = 0. \quad (\text{B.2})$$

Changing the names of indices we write (B.1) three times:

$$\begin{aligned} Q_{\lambda\kappa;\nu\mu} - Q_{\lambda\kappa;\mu\nu} &= Q_{\sigma\kappa} R^\sigma{}_{\lambda\nu\mu} + Q_{\lambda\sigma} R^\sigma{}_{\kappa\nu\mu}, \\ Q_{\mu\kappa;\lambda\nu} - Q_{\mu\kappa;\nu\lambda} &= Q_{\sigma\kappa} R^\sigma{}_{\mu\lambda\nu} + Q_{\mu\sigma} R^\sigma{}_{\kappa\lambda\nu}, \\ Q_{\nu\kappa;\mu\lambda} - Q_{\nu\kappa;\lambda\mu} &= Q_{\sigma\kappa} R^\sigma{}_{\nu\mu\lambda} + Q_{\nu\sigma} R^\sigma{}_{\kappa\mu\lambda}. \end{aligned}$$

We add the first equation, subtract the second one and finally add the third equation. Hence we obtain

$$\begin{aligned} & Q_{\lambda\kappa;\nu\mu} - Q_{\lambda\kappa;\mu\nu} - Q_{\mu\kappa;\lambda\nu} + Q_{\mu\kappa;\nu\lambda} + Q_{\nu\kappa;\mu\lambda} - Q_{\nu\kappa;\lambda\mu} \\ &= 2Q_{\lambda\kappa;\nu\mu} - (Q_{\lambda\kappa;\mu} + Q_{\mu\kappa;\lambda})_{;\nu} + (Q_{\mu\kappa;\nu} + Q_{\nu\kappa;\mu})_{;\lambda} - (Q_{\nu\kappa;\lambda} + Q_{\lambda\kappa;\nu})_{;\mu} \\ &= Q_{\sigma\lambda} R^\sigma{}_{\kappa\mu\nu} + Q_{\sigma\mu} R^\sigma{}_{\kappa\lambda\nu} + Q_{\sigma\nu} R^\sigma{}_{\kappa\lambda\mu} + 2Q_{\sigma\kappa} R^\sigma{}_{\mu\nu\lambda}. \end{aligned} \quad (\text{B.3})$$

In the last equality we used the standard property of curvature tensor: $R^\sigma_{[\mu\nu]\lambda} = 0$. Definition (2.1) applied to the terms in brackets in the above formula implies

$$\begin{aligned} 2Q_{\lambda\kappa;\nu\mu} &= \frac{2}{n-1} (g_{\lambda\mu}\xi_{\kappa;\nu} + g_{\nu\lambda}\xi_{\kappa;\mu} - g_{\mu\nu}\xi_{\kappa;\lambda} - g_{\kappa(\lambda}\xi_{\mu);\nu} + g_{\kappa(\mu}\xi_{\nu);\lambda} - g_{\kappa(\nu}\xi_{\lambda);\mu}) \\ &\quad + Q_{\sigma\lambda}R^\sigma_{\kappa\mu\nu} + Q_{\sigma\mu}R^\sigma_{\kappa\lambda\nu} + Q_{\sigma\nu}R^\sigma_{\kappa\lambda\mu} + 2Q_{\sigma\kappa}R^\sigma_{\mu\nu\lambda} \\ &\quad + \mathcal{Q}_{\lambda\kappa\mu;\nu} - \mathcal{Q}_{\mu\kappa\nu;\lambda} + \mathcal{Q}_{\nu\kappa\lambda;\mu}, \end{aligned} \quad (\text{B.4})$$

where the covector $\xi_\mu = \nabla^\lambda Q_{\lambda\mu}$ (cf. (2.6)). Hence for a CYK tensor Q we obtain (2.14).

Contracting (B.4) with respect to indices μ and ν and using the basic properties (2.2) of $\mathcal{Q}$ we get

$$Q_{\lambda\kappa}{}^{;\mu}{}_\mu + R^\sigma_{\kappa\lambda\mu}Q^\mu{}_\sigma + Q_{\sigma\kappa}R^\sigma_{\lambda} + \frac{2}{n-1}\xi_{(\kappa;\lambda)} + \frac{1}{n-1}g_{\kappa\lambda}\xi^\mu{}_{;\mu} = \nabla^\mu \mathcal{Q}_{\mu\kappa\lambda} - \frac{n-4}{n-1}\xi_{\kappa;\lambda}. \quad (\text{B.5})$$

Formulae (2.16) and (B.5) imply the following equation for a CYK tensor Q :

$$\nabla^\mu \nabla_\mu Q_{\lambda\kappa} = R^\sigma_{\kappa\lambda\mu}Q^\mu{}_\sigma - R_{\sigma[\kappa}Q_{\lambda]}{}^\sigma \quad (\text{B.6})$$

for $\dim M = 4$. It is interesting to point out that for compact four-dimensional Riemannian manifolds equation (B.6) is equivalent to Definition 1:

Theorem 9. *Let M be a compact (without boundary) Riemannian manifold with $\dim M = 4$. Then a two-form Q is a CYK tensor iff*

$$\nabla^\mu \nabla_\mu Q_{\lambda\kappa} = R^\sigma_{\kappa\lambda\mu}Q^\mu{}_\sigma + R_{\sigma[\lambda}Q_{\kappa]}{}^\sigma.$$

Proof. We need to show that equation (B.6) implies $\mathcal{Q}_{\lambda\kappa\mu}(Q, g) = 0$.

Similarly to (2.16) we derive

$$\frac{2}{3}\xi_{(\mu;\lambda)} + \frac{1}{3}g_{\mu\lambda}\xi^\nu{}_{;\nu} - R_{\sigma(\mu}Q_{\lambda)}{}^\sigma + \frac{1}{2}\nabla^\sigma \mathcal{Q}_{\lambda\sigma\mu} = 0, \quad 4\xi^\mu{}_{;\mu} + \nabla^\sigma \mathcal{Q}_{\nu\sigma\mu}g^{\mu\nu} = 0,$$

which together with (B.2) and (B.5) gives¹⁶

$$\nabla^\mu \nabla_\mu Q_{\lambda\kappa} + R^\sigma_{\kappa\lambda\mu}Q^\mu{}_\sigma + R_{\sigma[\kappa}Q_{\lambda]}{}^\sigma = \nabla^\mu \mathcal{Q}_{\mu\kappa\lambda} + \frac{1}{2}\nabla^\sigma \mathcal{Q}_{\kappa\sigma\lambda}. \quad (\text{B.7})$$

Contracting the above equality with Q and assuming equation (B.6) we get

$$\begin{aligned} 0 &= \left(\nabla^\mu \mathcal{Q}_{\mu\kappa\lambda} + \frac{1}{2}\nabla^\sigma \mathcal{Q}_{\lambda\sigma\kappa} \right) Q^{\kappa\lambda} = \nabla^\mu (\mathcal{Q}_{\mu\kappa\lambda}Q^{\kappa\lambda}) - \mathcal{Q}_{\mu\kappa\lambda}\nabla^\mu Q^{\kappa\lambda} \\ &= \nabla^\mu (\mathcal{Q}_{\mu\kappa\lambda}Q^{\kappa\lambda}) + \frac{1}{2}\mathcal{Q}_{\lambda\kappa\mu}Q^{\lambda\kappa\mu}. \end{aligned} \quad (\text{B.8})$$

Finally, we integrate the above formula over M , a total divergence drops out, and the integral $\int_M \sqrt{\det g} \mathcal{Q}_{\lambda\kappa\mu}Q^{\lambda\kappa\mu}$ vanishes. This implies $\mathcal{Q}^{\lambda\kappa\mu} = 0$. $\square$

A similar result holds for a p -form Q in $2p$ -dimensional M (see [38]).

¹⁶For a CYK tensor Q formula (B.7) obviously gives equation (B.6).

B.1 Miscellaneous results

It is worth pointing out that equation (B.6) can be derived¹⁷ from a variational principle $\delta \int_M \mathcal{L} = 0$, where $\mathcal{L}[Q] = \frac{1}{4} \sqrt{\det g} \mathcal{Q}_{\lambda\kappa\mu} \mathcal{Q}^{\lambda\kappa\mu}(Q, g)$.

Equation (B.6) for Ricci flat metrics gives

$$2\nabla^\mu \nabla_\mu Q_{\kappa\lambda} = 2R^\sigma{}_{\kappa\lambda\mu} Q^\mu{}_\sigma = R_{\kappa\lambda\sigma\mu} Q^{\sigma\mu} = F_{\kappa\lambda}(W, Q)$$

and, in particular, it is true for $Q = Y$ in Kerr spacetime. Moreover,

$$\frac{1}{4} F_{\mu\nu}(W, Y) Y^{\mu\nu} = \frac{2mr}{\rho^2} = \frac{2mr}{r^2 + a^2 \cos^2 \theta} = g(\partial_t, \partial_t) + 1$$

gives a nice density in Boyer-Lindquist coordinates:

$$\sqrt{-g} F_{\mu\nu}(W, Y) Y^{\mu\nu} = 8mr \sin \theta.$$

A complex-valued function $\zeta := r + \imath \cos \theta$ enables one to write some objects in Kerr spacetime in a compact form like the Ernst potential $\mathcal{E} = 1 - 2m/\bar{\zeta}$ or Newman–Penrose scalar $\Psi_2 = -m/\bar{\zeta}^3$ (cf. [40]), where by $\bar{\zeta}$ we denote the complex conjugate i.e. $\bar{\zeta} := r - \imath \cos \theta$. In this context in Eddington–Finkelstein coordinates we have

$$Y + \imath * Y = -\imath \zeta d\zeta \wedge (dv - a \sin^2 \theta d\tilde{\phi}) + \bar{\zeta} \zeta^2 \sin \theta d\theta \wedge d\tilde{\phi},$$

$$\begin{aligned} K' + \imath K'' &= \frac{\bar{\zeta} - \zeta}{2\zeta} \left(a^2 \sin^2 \theta \partial_v \otimes \partial_v + \partial_\theta \otimes \partial_\theta + \frac{1}{\sin^2 \theta} \partial_{\tilde{\phi}} \otimes \partial_{\tilde{\phi}} + 2a \partial_v \otimes_s \partial_{\tilde{\phi}} \right) \\ &\quad + \frac{\bar{\zeta} + \zeta}{2\zeta} (\Delta \partial_r \otimes \partial_r + 2(r^2 + a^2) \partial_v \otimes_s \partial_r + 2a \partial_r \otimes_s \partial_{\tilde{\phi}}), \end{aligned}$$

and, in particular, $\text{tr}(K' + \imath K'') = 2\bar{\zeta}^2 = \frac{\mathcal{E}-1}{\Psi_2}$.

References

- [1] L.F. Abbott and S. Deser, *Stability Of Gravity With A Cosmological Constant*, Nucl. Phys. B 195, (1982) 76–96
- [2] L. Andersson, *Bel–Robinson Energy and Constant Mean Curvature Foliations*, Ann. Henri Poincaré 5, (2004) 235–244
- [3] S. Benenti and M. Francaviglia, *General Relativity and Gravitation* vol. 1, ed. A. Held (New York: Plenum), (1980) 393–439
- [4] I.M. Benn, P. Charlton and J. Kress, *Debye potentials for Maxwell and Dirac fields from a generalization of the Killing–Yano equation*, Journal of Mathematical Physics 38, (1997) 4504–4527

¹⁷The calculations are similar to the proof of Theorem 9.

- [5] I.M. Benn and P. Charlton, *Dirac symmetry operators from conformal Killing-Yano tensors*, Classical and Quantum Gravity 14, (1997) 1037–1042
- [6] G. Bergqvist, P. Lankinen, *Unique characterization of the Bel–Robinson tensor*, Classical and Quantum Gravity 21, (2004) 3489–3503; G. Bergqvist, I. Eriksson and J.M.M. Senovilla, *New electromagnetic conservation laws*, Classical and Quantum Gravity 20, (2003) 2663–68
- [7] M.A.G. Bonilla and J.M.M. Senovilla, *Some Properties of the Bel and Bel–Robinson Tensors*, General Relativity and Gravitation 29, (1997) 91–116
- [8] H.L. Bray, P.T. Chruściel, *The Penrose Inequality*, in: The Einstein Equations and the Large Scale Behavior of Gravitational Fields (50 Years of the Cauchy Problem in General Relativity), P.T. Chruściel, H. Friedrich, Eds, Birkhaeuser, Basel, 2004, pp. 39–70, <http://arxiv.org/abs/gr-qc/0312047>
- [9] R. Casalbuoni, *Relativity And Supersymmetries*, Phys. Lett. B **62** (1976) 49
- [10] B. Carter, *Global structure of the Kerr family of gravitational fields*, Phys. Rev. 174, (1968) 1559–1571; *Hamilton–Jacobi and Schrödinger separable solutions of Einstein’s equations*, Commun. Math. Phys. 10, (1968) 280; *Separability Of The Killing Maxwell System Underlying The Generalized Angular Momentum Constant In The Kerr–Newman Black Hole Metrics*, Journal of Mathematical Physics 28, (1987) 1535–1538
- [11] D. Christodoulou, *Reversible and irreversible transformations in black hole physics*, Phys. Rev. Letters 25, (1970) 1596–1597; D. Christodoulou and R. Ruffini, *Reversible transformations of a charged black hole*, Phys. Rev. D 4, (1971) 3552–3555
- [12] D. Christodoulou and S. Klainerman, *Asymptotic Properties of Linear Field Equations in Minkowski Space*, Communications on Pure and Applied Mathematics 43, (1990) 137–199
- [13] C.D. Collinson and L. Howarth, *Generalized Killing Tensors*, General Relativity and Gravitation 32, (2000) 1767–1776
- [14] S.R. Douglas, *Letter: Review of the Definitions of the Bel and Bel–Robinson Tensors*, General Relativity and Gravitation 35, (2003) 1691–97
- [15] F. Fayos and C.F. Sopuerta, *On the Papapetrou field in vacuum*, Classical and Quantum Gravity 16, (1999) 2965–85
- [16] F. de Felice, *Angular momentum and separation constant in the Kerr metric*, J. Phys. A: Math. Gen. 13, (1980) 1701–1708
- [17] V.P. Frolov and I.D. Novikov, *Black Hole Physics: Basic Concepts and New Developments*, Kluwer Academic Pub, (1998); in Russian: Новиков И. Д., Фролов В. П., *Физика черных дыр*, Наука, (Москва 1986)

- [18] G.W. Gibbons and P.J. Ruback, *The Hidden Symmetries of Multi-Centre Metrics*, Commun. Math. Phys. 115, (1988) 267–300
- [19] G.W. Gibbons, R.H. Rietdijk and J.W. van Holten, *SUSY in the sky*, Nucl. Phys. B 404, (1993) 42–64, <http://arxiv.org/abs/hep-th/9303112>
- [20] E.N. Glass and J. Kress, *Solutions of Penrose’s Equation*, Journal of Mathematical Physics 40, (1999) 309–317, <http://arxiv.org/abs/gr-qc/9809074>
- [21] E.N. Glass and M.G. Naber, *Gravitational mass anomaly*, Journal of Mathematical Physics 35, (1994) 4178–83
- [22] J.N. Goldberg, *Conserved quantities at spatial and null infinity: The Penrose potential*, Phys. Rev. D 41, (1990) 410–417
- [23] M. Heusler, *Black Hole Uniqueness Theorems*, Cambridge Lecture Notes in Physics 6, University Press, p. 119 (Cambridge 1996)
- [24] L.P. Hughston and P. Sommers, *Spacetimes with Killing Tensors*, Commun. math. Phys. 32, (1973) 147–152
- [25] J. Jezierski, *The Relation between Metric and Spin-2 Formulations of Linearized Einstein Theory*, Gen. Rel. and Grav. 27, (1995) 821–843, <http://arxiv.org/abs/gr-qc/9411066>
- [26] J. Jezierski, *Conformal Yano-Killing tensors and asymptotic CYK tensors for the Schwarzschild metric*, Classical and Quantum Gravity 14, (1997) 1679–1688, <http://arxiv.org/abs/hep-th/9411074>
- [27] J. Jezierski, *Gauge-invariant formulation of the linearized Einstein equations on the Schwarzschild background*, in Current Topics in Math. Cosmology, eds M. Rainer and H-J Schmidt, World Scientific (1998); *Energy and angular momentum of the weak gravitational waves on the Schwarzschild background – quasilocal gauge-invariant formulation*, General Relativity and Gravitation 31, (1999) 1855–1890, <http://arxiv.org/abs/gr-qc/9801068>
- [28] J. Jezierski, *CYK Tensors, Maxwell Field and Conserved Quantities for Spin-2 Field*, Classical and Quantum Gravity 19, (2002) 4405–4429, <http://arxiv.org/abs/gr-qc/0211039>
- [29] D. Kastor and J. Traschen, *Conserved gravitational charges from Yano tensor*, <http://arxiv.org/abs/hep-th/0406052>
- [30] A. Moroianu, U. Semmelmann, *Twistor forms on Kähler manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) II (2003), 823–845, <http://arxiv.org/abs/math.DG/0204322>
- [31] A. Papapetrou, Ann. Inst. H Poincaré A 4, (1966) 83

- [32] N. Pelavas, N. Neary and K. Lake, *Properties of the instantaneous Ergo Surface of a Kerr Black Hole*, <http://arxiv.org/abs/gr-qc/0012052>
- [33] R. Penrose, *Gravitational Collapse: The Role of General Relativity*, Riv. del Nuovo Cimento 1, (1969) 252–276
- [34] R. Penrose, *Naked Singularities*, Ann. N.Y. Acad. Sci. 224, (1973) 125–134; R. Floyd, *The Dynamics of Kerr Fields*, PhD. Thesis (London 1973); B. Carter and R.G. McLenaghan, *Generalized Total Angular Momentum Operator for The Dirac Equation in Curved Space-Time*, Phys. Rev. D 19, (1979) 1093–1097
- [35] R. Penrose, *Quasi-local Mass and Angular Momentum in General Relativity*, Proc. Roy. Soc. Lond. A381, (1982) 53–62
- [36] R. Penrose and W. Rindler, *Spinors and Space-time*, Cambridge University Press, Vol. 2, p. 396 (Cambridge 1986)
- [37] R. Rani, S.B. Edgar and A. Barnes, *Killing tensors and conformal Killing tensors from conformal Killing vectors*, Classical and Quantum Gravity 20, (2003) 1929–1942
- [38] U. Semmelmann, *Conformal Killing forms on Riemannian manifolds*, Mathematische Zeitschrift 245, (2003) 503–527
- [39] S.E. Stepanov, *The Vector Space of Conformal Killing Forms on a Riemannian Manifold*, Journal of Mathematical Sciences 110, (2002) 2892–2906; *On conformal Killing 2-form of the electromagnetic field*, Journal of Geometry and Physics 33, (2000) 191–209
- [40] H. Stephani et al., *Exact solutions of Einstein’s field equations*, 2nd ed., University Press (Cambridge 2003)
- [41] M. Walker and R. Penrose, *On quadratic first integrals of the geodesic equations for type {22} spacetimes*, Commun. Math. Phys. 18, (1970) 265–274
- [42] S. Tachibana, *On conformal Killing tensor in a Riemannian space*, Tohoku Math. J. (2) 21, (1969) 56–64; T. Kashiwada, *On conformal Killing tensor*, Natur. Sci. Rep. Ochanomizu Univ. 19, (1968) 67–74; S. Tachibana and T. Kashiwada, *On the integrability of Killing–Yano’s equation*, J. Math. Soc. Japan 21, (1969) 259–265
- [43] N.M.J. Woodhouse, *Killing Tensors and the Separation of the Hamilton–Jacobi Equation*, Commun. Math. Phys. 44, (1975) 9–38
- [44] K. Yano, *Some remarks on tensor fields and curvature*, Ann. Math. 55, (1952) 328–347